

SUPPLEMENTARY TABLE S7. SEX-SPECIFIC ASSOCIATIONS (β [CI]) BETWEEN BDE-47 TRAJECTORIES (NG/G LIPID) AND SERUM THYROID PARAMETERS

	<i>High decreasing vs. persistent low</i>			<i>Low increasing vs. persistent low</i>			<i>p-int</i>
	<i>Boys</i>	<i>Girls</i>	<i>p-int</i>	<i>Boys</i>	<i>Girls</i>	<i>p-int</i>	
Log ₁₀ TSH (μ IU/mL)	-0.06 [-0.15 to 0.03]	-0.10 [-0.20 to -0.00]	0.54	-0.15 [-0.26 to -0.05]	-0.04 [-0.13 to 0.05]	0.12	
Free T4 (pmol/L)	-0.27 [-1.32 to 0.79]	-0.12 [-1.06 to 0.83]	0.78	-1.19 [-2.18 to -0.20]	-0.19 to [-1.19 to 0.82]	0.21	
Total T4 (nmol/L)	-4.50 [-21.69 to 12.69]	3.15 [-7.94 to 14.23]	0.67	-14.28 [-27.48 to -1.07]	2.80 [-7.32 to 12.92]	0.06	

185 children and 223 observations.