

**Table S2.** Risk of mortality on dialysis associated with HCV serostatus in multivariable Cox regression (N=414,761)

	<b>HR</b>	<b>95% CI</b>	<b>P value</b>
<b>HCV-seronegative</b>	ref		
<b>HCV-seropositive</b>	1.09	1.07-1.11	<0.001
<b>Age &lt;35 years</b>			
<b>35-45</b>	1.37	1.32-1.42	<0.001
<b>45-55</b>	1.91	1.85-1.98	<0.001
<b>55-65</b>	2.68	2.59-2.77	<0.001
<b>65-75</b>	3.77	3.65-3.91	<0.001
<b>&gt;75</b>	5.69	5.50-5.89	<0.001
<b>Male</b>	1.03	1.01-1.03	<0.001
<b>Race</b>			
<b>Caucasian</b>	ref		
<b>African American</b>	0.65	0.65-0.66	<0.001
<b>Asian</b>	0.55	0.54-0.57	<0.001
<b>Latino</b>	0.56	0.56-0.58	<0.001
<b>Other</b>	0.67	0.66-0.69	<0.001
<b>Preobservation dialysis time</b>	1.02	1.02-1.02	<0.001
<b>Hemodialysis</b>	ref		
<b>Peritoneal dialysis</b>	0.83	0.81-0.85	<0.001
<b>Medicare</b>	ref		
<b>Medicaid</b>	1.16	1.14-1.20	<0.001
<b>Private</b>	0.92	0.91-0.93	<0.001
<b>Income quintile (US\$)</b>			
<b>\$8,993-35,363</b>	ref		
<b>\$35,370-43,265</b>	1.00	0.99-1.02	0.236
<b>\$43,331-52,143</b>	1.01	0.99-1.02	0.200
<b>\$52,173-67,316</b>	0.94	0.93-0.95	<0.001
<b>\$67,331-180,120</b>	0.91	0.89-0.92	<0.001
<b>CVD</b>	1.04	1.02-1.05	<0.001
<b>COPD</b>	1.24	1.21-1.27	<0.001
<b>Diabetes mellitus</b>	1.11	1.09-1.12	<0.001
<b>Liver disease</b>	1.17	1.12-1.21	<0.001
<b>Albumin ≥3.5 (g/dL)</b>	ref		
<b>&lt;3.5</b>	1.62	1.61-1.64	<0.001
<b>Hemoglobin ≥12 (g/dL)</b>	ref		
<b>10-12</b>	1.05	1.04-1.06	<0.001
<b>&lt;10</b>	1.06	1.04-1.07	<0.001
<b>Platelets ≥150 x 10<sup>3</sup>/mm<sup>3</sup></b>	Ref		
<b>&lt;150</b>	1.40	1.38-1.42	<0.001

**Abbreviations:** HR – Hazard ratio; CI – Confidence Interval; HCV – Hepatitis C virus; CVD – cardiovascular disease; COPD – chronic obstructive pulmonary disease; BMI – body mass index; AVF – arteriovenous fistula; AVG – arteriovenous graft; CVC – central venous catheter; PD – peritoneal dialysis; FIB-4 – Fibrosis-4 Index