

Suppl. table 1. Associations of FLI with free and protein-bound AGEs and sRAGE

FLI					
Outcome	Model	β	95% CI		P-value
Free CML (nM)	1	0.031	0.001	0.062	0.043
	2	0.027	-0.004	0.057	0.083
	3	0.033	0.002	0.064	0.038
	4	0.048	-0.004	0.099	0.068
Free CEL (nM)	1	0.064	0.034	0.093	<0.001
	2	0.053	0.023	0.083	<0.001
	3	0.058	0.029	0.088	<0.001
	4	0.069	0.019	0.119	0.007
Free MG-H1 (nM)	1	0.013	-0.034	0.060	0.590
	2	0.000	-0.047	0.047	0.990
	3	-0.001	-0.048	0.047	0.976
	4	0.004	-0.076	0.083	0.930
PB-Pentosidine (nmol/mmol lysine)	1	-0.070	-0.101	-0.039	<0.001
	2	-0.086	-0.117	-0.055	<0.001
	3	-0.093	-0.124	-0.062	<0.001
	4	-0.061	-0.113	-0.009	0.022
PB-CML (nmol/mmol lysine)	1	-0.089	-0.110	-0.067	<0.001
	2	-0.096	-0.117	-0.074	<0.001
	3	-0.081	-0.103	-0.058	<0.001
	4	-0.063	-0.100	-0.025	0.001
PB-CEL (nmol/mmol lysine)	1	0.026	-0.002	0.053	0.067
	2	0.024	-0.005	0.003	0.621
	3	0.033	0.003	0.063	0.031
	4	0.046	-0.004	0.096	0.069
sRAGE (pg/ml)	1	-0.090	-0.127	-0.053	<0.001
	2	-0.066	-0.103	-0.029	0.001
	3	-0.063	-0.102	-0.025	0.001
	4	-0.030	-0.095	0.034	0.354

Suppl. table 2. Associations of FLI with low-grade inflammation

FLI					
Outcome	Model	β	95% CI		P-value
LGI	1	0.325	0.242	0.408	<0.001
	2	0.343	0.260	0.427	<0.001
	3	0.329	0.243	0.416	<0.001
	4	0.234	0.090	0.378	0.002