

Supplementary Table S1. Primers used for qRT-PCR in this study

<b>Gene</b>	<b>Forward (5' -&gt; 3')</b>	<b>Reverse (5' -&gt; 3')</b>
<i>CCA1</i>	TCTGGTTATTAAGACTCGGAAGCCAT	CTTCTGCCATGCTCTACCATAAAGCC
<i>PRR9</i>	CCTTGAAGATACTGATGAAACTTGTG	TGTAATTGATAGCCTGAGCACCCTT
<i>PRR7</i>	ACGACTGAGAACAACGCTTTCACAAA	ACATGGACAAGGTTATAGGAGGCATG
<i>GI</i>	AATTCAGCACGCGCCTATTG	GTTGCTTCTGCTGCAGGAACTT
<i>PIF4</i>	ACCTCAGAGACGGTTAAGCC	TGGAGGAGGCATGACTTGAG
<i>U-box</i>	TGCGCTGCCAGATAATACTACTATT	TGCTGCCCAACATCAGGTT

Primers used for plasmid construction in this study

<b>Gene</b>	<b>Forward (5' -&gt; 3')</b>	<b>Reverse (5' -&gt; 3')</b>
<i>PRR7pro</i>	ATCGATATCCCTTTTGCAGTCACGATGAT	GTAGATATCCACACCAACTCTGCTTCGCT
<i>PRR7</i>	GCTACCATGGATGAATGCTAATGAGGAG GG	TAGCCCATGGCGCTATCCTCAATGTTTTTAA
<i>TAP-ELF3</i>	CACCGCATGAAGAGAGGGAAAGATGAG G	TTAAGGCTTAGAGGAGTCA
<i>HA-ELF3</i>	CACCATGAAGAGAGGGAAAGAT	TTAAGGCTTAGAGGAGTCA
<i>ELF3<sup>A37T</sup></i>	AGAAACAAGATGACTCTTTATGAGCAG	CTGCTCATAAAGAGTCATCTTGTTTCT
<i>ELF3<sup>P666S</sup></i>	ATAAAGGTGGTATCTCACAAACGCAAAG	CTTTGCGTTGTGAGATACCACCTTTAT
<i>ELF4</i>	CACCATGAAGAGGAACGGCGAG	TTAAGCTCTAGTTCCGGC
<i>LUX</i>	CACCATGGGAGAGGAAGTACAAA	ATTCTCATTTGCGCTTCC
<i>GI</i>	CACCATGGCTAGTTCATCTTCA	TTATTGGGACAAGGATATAG
<i>PIF4</i>	CACCATGGAACACCAAGGTTGGAG	CTAGTGGTCCAACGAGAACCG