

**Supplementary Table S2: Demographic and health information of prostate cancer patients.**

		<b>Non-Treated (n=41)</b>	<b>Treated with Abiraterone Only (n=16)</b>	<b>Treated with Enzalutamide Only (n=13)</b>
<b>Race</b>	<i>Caucasian</i>	30 (73.2%)	9 (56.3%)	11 (84.6%)
	<i>African American</i>	10 (24.4%)	7 (43.7%)	2 (15.4%)
	<i>Unknown</i>	1 (2.4%)	0 (0%)	0 (0%)
<b>Age</b>	<i>18-64</i>	18 (43.9%)	3 (18.8%)	3 (23.1%)
	<i>≥65</i>	23 (56.1%)	13 (81.2%)	10 (76.9%)
<b>Metastasis</b>	<i>None</i>	16 (39.0%)	1 (6.3%)	0 (0%)
	<i>Bone</i>	23 (56.1%)	14 (87.4%)	11 (84.6%)
	<i>Other Sites</i>	2 (4.9%)	1 (6.3%)	2 (15.4%)
<b>Gleason Score</b>	<i>&lt;4</i>	0 (0%)	0 (0%)	0 (0%)
	<i>&gt;4</i>	38 (92.7%)	14 (87.5%)	13 (100%)
	<i>Unknown</i>	3 (7.3%)	2 (12.5%)	0 (0%)
<b>PSA Level Post-Therapy (ng/mL)</b>	<i>&lt;4</i>	24 (58.5%)	7 (43.7%)	5 (38.5%)
	<i>&gt;4</i>	17 (41.5%)	9 (56.3%)	8 (61.5%)
	<i>Unknown</i>	0 (0%)	0 (0%)	0 (0%)
<b>Family History of Cancer</b>	<i>No</i>	12 (29.3%)	5 (31.3%)	3 (23.1%)
	<i>Yes</i>	25 (61%)	10 (62.4%)	8 (61.5%)
	<i>Unknown</i>	4 (9.7%)	1 (6.3%)	2 (15.4%)
<b>Prostatectomy</b>	<i>No</i>	0 (0%)	8 (50%)	4 (30.8%)
	<i>Yes</i>	15 (36.6%)	8 (50%)	6 (46.2%)
	<i>Unknown</i>	26 (63.4%)	0 (0%)	3 (23%)
<b>Tobacco Use</b>	<i>No</i>	13 (31.7%)	4 (25%)	5 (38.5%)
	<i>Yes</i>	27 (65.9%)	12 (75%)	8 (61.5%)
	<i>Unknown</i>	1 (2.4%)	0 (0%)	0 (0%)