

Supplementary Table 1. Multivariable-adjusted HRs (95% CIs) for incident ischemic stroke according to family history of stroke

Variable	Family history of stroke	
	No	Yes
Total population		
No. of participants	401,227	85,971
No. of ischemic stroke cases	28,556	9,023
Cases per 1,000 PYs	7.33	10.89
HR (95% CI)		
Sex adjusted	Ref	1.25 (1.22–1.28)
Multivariable adjusted*	Ref	1.19 (1.16–1.22)
Men		
No. of participants	163,791	35,449
No. of ischemic stroke cases	12,933	3,990
Cases per 1,000 PYs	8.30	11.89
HR (95% CI)		
Crude	Ref	1.25 (1.21–1.30)
Multivariable adjusted*	Ref	1.19 (1.15–1.23)
Women		
No. of participants	237,436	50,522
No. of ischemic stroke cases	15,623	5,033
Cases per 1,000 PYs	6.69	10.21
HR (95% CI)		
Crude	Ref	1.25 (1.21–1.29)
Multivariable adjusted*	Ref	1.19 (1.15–1.23)

Multivariable model was stratified by age and study area.

HR, hazard ratio; CI, confidence interval; PY, person-year.

*Adjusted for level of education; marital status; household income; alcohol consumption; smoking status; physical activity; intake frequencies of red meat, fresh fruits, and vegetables; menopause status (only for women); body mass index; and prevalent diabetes and hypertension.