	Prog	rip	Suture fix	ation		Odds Ratio		Odds Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% CI		M-H, Random, 95% Cl
Bruna Estaban 2010	2	38	4	39	6.6%	0.49 [0.08, 2.83]		
Chatzimavroudis 2014	1	25	1	25	2.6%	1.00 [0.06, 16.93]		
Jorgensen 2013	16	161	13	168	34.9%	1.32 [0.61, 2.83]		
Molegraaf 2017	9	164	8	166	21.4%	1.15 [0.43, 3.05]		_
Ronka 2015	15	189	8	197	26.3%	2.04 [0.84, 4.92]		
Verhagen 2016	7	161	2	165	8.1%	3.70 [0.76, 18.11]		
Total (95% CI)		738		760	100.0%	1.45 [0.92, 2.28]		-
Total events	50		36					
Heterogeneity: Tau ² = 0.00; Chi ² = 3.75, df = 5 (P = 0.59); $I^2 = 0\%$							0.05	0.2 1 5 20
Test for overall effect: $Z = 1.61$ (P = 0.11)							0.05	Favours [Progrip] Favours [Suture fixation]

Supporting information: figure 6 Forest plot comparing the incidence of chronic pain (VAS>3) of $ProGrip^{TM}$ and suture fixation at 12 months postoperatively.