

**Supplementary Table 1.** Characteristics of cohort for prospective study of menopause and incident kidney stones by menopausal status at study midpoint, 2003

	Pre-menopausal (N=52,503)	Post-menopausal (N=31,522)
Age (years), mean (SD)	46 (4)	52 (4)
Race, white	48,661 (93%)	29,059 (92%)
Body mass index (kg/m <sup>2</sup> )	26.6 (6.2)	27.6 (6.5)
Menopause etiology		
Natural		20,443 (65%)
Surgical		10,139 (32%)
Radiation		940 (3%)
Post-menopausal hormone therapy use		
Never		9,846 (31%)
Past		9,642 (31%)
Current		11,517 (37%)
History of hypertension	8,265 (16%)	9,225 (29%)
History of diabetes	1,297 (2.5%)	1,641 (5.2%)
Thiazide use	1,881 (3.6%)	2,250 (7.1%)
Fluid intake (L/day)	1.7 (0.9)	1.7 (1.0)
Supplemental calcium (mg/d), median (25%, 75%)	200 (0, 662)	500 (23, 1000)
Total vitamin D intake (IU)	452 (305)	545 (343)

Abbreviations: SD, standard deviation

\*Data are presented as % unless otherwise indicated. Dietary intake factors are presented as mean and standard deviation (SD) unless otherwise indicated.

