

Supplementary

Life History Effects Linked to an Advantage for *wAu* *Wolbachia* in *Drosophila*

Li-Jun Cao ^{1,2,*}, Weibin Jiang ^{1,3} and Ary A. Hoffmann ^{1,*}

¹ Pest and Adaptation Research Group, School of BioSciences, Bio21 Institute, The University of Melbourne, Melbourne, VIC 3010, Australia

² Institute of Plant and Environmental Protection, Beijing Academy of Agriculture and Forestry Sciences, Beijing 100097, China

³ College of Life & Environmental Science, Shanghai Normal University, Shanghai 200234, China; jiangwb@shnu.edu.cn

* Correspondence: gmatjhpl@163.com (L.-J.C.); ary@unimelb.edu.au (A.A.H.)

Received: 8 April 2019; Accepted: 30 April 2019; Published: date



Figure S1. Fungus-infestation levels on grapes with three levels recognized: (a) Level 1, less than 1 piece of 6 pieces of grapes was covered by visible mycelial growth; (b) level 2, 2 to 4 pieces of grapes were covered by visible growth; (c) level 3, 5 to 6 pieces of grapes were covered by visible growth.



© 2019 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).