
1	Flow Density in Optical Coherence Tomography
2	Angiography is Useful for Retinopathy Diagnosis in
3	Diabetic Patients
4 5 6	Yoshihiro Kaizu, ¹ Shintaro Nakao, ¹ Mitsuru Arima, ¹ Takehito Hayami, ² Iori Wada, ¹ Muneo Yamaguchi, ¹ Haruka Sekiryu, ¹ Keijiro Ishikawa, ¹ Yasuhiro Ikeda, ¹ Koh-Hei Sonoda ¹
7 8	Address: ¹ Department of Ophthalmology, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan.
9 10	² Department of Intelligent Mechanical Systems, Graduate School of Natural Science and Technology, Okayama University, Okayama, Japan
11 12 13	Short Title: Flow Density in OCTA for DR diagnosis
14	Correspondence:
15	Shintaro Nakao, MD, PhD
16	Kyushu University
17	3-1-1 Maidashi, Higashi-Ku, Fukuoka 812-8582, Japan.
18	Tel: 81-92-642-5648
19	Fax: 81-92-642-5663
20	Email: <u>snakao@med.kyushu-u.ac.jp</u>



Supplemental Fig. 1. Quantification of flow density in OCTA. The OCTA image of a 45-year-old man at the SCP. (A) A 3 X 3 mm en face OCTA was cropped to 2.6 X 2.6 mm using an image editing program (Adobe Systems, Inc.; yellow dotted square) to avoid the influence of the "angioFLOW" display on quantification. (B) A 2.6 X 2.6 mm en face OCTA image used for this study. (C) Image binarization was performed using NI Vision. (D) Each image was uniformly divided into four regions (superior, inferior, nasal and temporal sides).