

**Table A. Attrition analysis, n=3133, North Carolina Youth Tobacco Survey, 2017 <sup>a</sup>**

	Included in final sample, n=2835 (89.5%) n (%)	Not included in final sample, n=298 (10.5%) n (%)	P-value
Sex			
Female	1428 (94.1%)	91 (5.9%)	p<0.001
Male	1407 (85.5%)	201 (14.5)	
Grade			
9 <sup>th</sup>	904 (92.8%)	62 (7.2%)	p=0.15
10 <sup>th</sup>	696 (91.0%)	67 (9.0%)	
11 <sup>th</sup>	577 (89.9%)	64 (10.1%)	
12 <sup>th</sup>	658 (87.7%)	79 (12.4%)	
Race			
Non-Hispanic White	1503 (91.8%)	112 (8.2%)	p=0.006
Non-Hispanic Black	650 (89.1%)	74 (10.9%)	
Hispanic	517 (86.6%)	77 (13.4%)	
Non-Hispanic other race	165 (84.9%)	23 (15.1%)	
Free or reduced-price lunch			
Yes	1362 (88.6%)	146 (11.4%)	p=0.13
No	1473 (91.1%)	131 (8.9%)	
Current cigarette use			
No	2617 (93.3%)	167 (6.7%)	p<0.001
Yes	218 (75.5%)	48 (24.5%)	
Current cigar use			
No	2497 (93.1%)	151 (6.9%)	p<0.001
Yes	338 (80.0%)	75 (20.0%)	
Current smokeless tobacco use			
No	2649 (92.9%)	188 (7.1%)	p<0.001
Yes	186 (67.0%)	71 (33.0%)	
Current waterpipe use			
No	2722 (93.1%)	179 (6.9%)	p<0.001
Yes	113 (68.3%)	35 (31.7%)	
Current tobacco e-cigarette use			
No	2334 (92.7%)	156 (7.3%)	p=0.004
Yes	501 (85.9%)	74 (14.1%)	

<sup>a</sup> Percentages provided are row percentages, i.e., the percentage in the cell included in the first column and first row indicates that among females, 94.1% were included in the final sample.