

Supplementary file 5

eTable 5. Distribution of missing and complete data across the study variables, the Stockholm Public Health Cohort			
Study variables	Missing data¹	Complete data²	P-value³
N (%)	4633 (23.5)	15087 (76.5)	
Age, mean (SD)	61.6 (13.1)	57.2 (11.2)	<0.001
Gender			<0.01
Men	44.9	46.7	
Women	55.1	53.3	
Country of origin			<0.001
Sweden	72.7	84.5	
Nordic	9.6	7.3	
Other	17.7	8.2	
Marital status			
Married	53.7	60.1	
Single/divorced/widowed	46.3	39.9	
Father's SEP			<0.001
Nonmanual	51.9	53.9	
Manual	48.1	46.1	
Educational attainment			<0.001
High	22.1	36.8	
Medium	33.7	38.9	
Low	44.2	24.3	
Adult SEP			<0.001
Nonmanual	53.0	64.4	
Manual	40.6	27.0	
Other	6.4	8.6	
Smoking			<0.001
Never smokers	50.3	46.8	
Current smokers	20.2	17.6	
Former smokers	29.5	35.6	
Risky alcohol drinking			<0.001
No	66.1	78.4	
Yes	33.9	21.6	
Physical inactivity			<0.001
Active	10.1	11.4	
Moderately active	15.8	21.3	
Slightly active	51.4	52.4	
Inactive	22.7	14.9	
Diet (Fruits and berries)			<0.001
More than once a day	19.0	22.9	
Almost daily/a few times a week	66.1	64.4	
Once a week or less	14.9	12.7	
Body Mass Index			<0.001
Underweight	2.0	1.1	
Normal weight	46.7	49.8	
Overweight	38.5	38.6	
Obese	12.8	10.5	

¹The sample with data missing on any of the study variables. ²The sample with complete data on all study variables. ³T-test for age and chi-square test for all categorical variables
