	Adjusted risk ratio	95% CI
Race		
Non-Hispanic white	1.00	Reference
Non-Hispanic black	1.20	1.14, 1.26
Hispanic	0.93	0.88, 0.99
Asian or Pacific Islander	1.06	0.95, 1.18
Native American	1.09	0.90, 1.33
Other	1.05	0.95, 1.16
Unknown	1.04	0.96, 1.13
Year		
2012	1.00	Reference
2013	1.04	0.99, 1.09
2014	1.04	0.99, 1.09
Hospital bed size		·
Small	1.00	Reference
Medium	1.03	0.96, 1.10
Large	1.07	1.00, 1.13
Insurance status		,
Private	1.00	Reference
Medicare	1.01	0.91, 1.12
Medicaid	0.95	0.91, 0.99
Other	0.99	0.90, 1.10
Uninsured	0.83	0.74, 0.92
Hospital Location		
Urban non-teaching	1.00	Reference
Rural	1.00	0.93, 1.09
Urban teaching	1.00	0.96, 1.05
Income Quartile		,
1 st (lowest)	1.00	Reference
2nd	1.00	0.95, 1.05
3rd	0.98	0.93, 1.03
4 th (highest)	0.95	0.89, 1.01
Hospital Region	0.55	0.00, 1.01
Northeast	1.00	Reference
Midwest	1.21	1.13, 1.29
South	1.27	1.19, 1.34
West	1.26	1.17, 1.35
Comorbidity index	1.20	1.17, 1.35
0	1.00	Reference
1-2	1.30	1.24, 1.35
>2	3.14	2.98, 3.31

Supplemental Table 1. Adjusted models for severe morbidity during postpartum readmissions with comorbidity index inclusive of maternal age

Adjusted model included all factors in this table (race, year, bed size, insurance status, hospital location, income quartile, hospital region, and comorbidity index score).