### Online Appendix

# A General Age-Specific Mortality Model with An Example Indexed by Child or Both Child and Adult Mortality

Samuel J. Clark 1,2,\*

<sup>1</sup>Department of Sociology, The Ohio State University

<sup>2</sup>MRC/Wits Rural Public Health and Health Transitions Research Unit (Agincourt), School of Public Health, Faculty of Health Sciences, University of the Witwatersrand

\*Contact: work@samclark.net, 206.303.9620

2018

#### Appendix A SVD Algebra

Below I rearrange the basic SVD relationship to derive useful additional relationships. **X** is an  $A \times L$  matrix of rank  $\rho$ .

$$\begin{array}{c|cccc}
\mathbf{X} = \mathbf{USV}^{\mathsf{T}} & (A.1) \\
\begin{bmatrix} \downarrow & & \downarrow \\ \mathbf{x}_{1} & \dots & \mathbf{x}_{L} \\ \downarrow & & \downarrow \end{bmatrix} = \begin{bmatrix} \downarrow & & \downarrow \\ \mathbf{u}_{1} & \dots & \mathbf{u}_{\rho} \\ \downarrow & & \downarrow \end{bmatrix} \begin{bmatrix} s_{1} & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & s_{\rho} \end{bmatrix} \begin{bmatrix} -\mathbf{v}_{1} & -\mathbf{v}_{\rho} \\ \vdots \\ -\mathbf{v}_{\rho} & -\mathbf{v}_{\rho} \end{bmatrix} \\
&= \begin{bmatrix} \downarrow & & \downarrow \\ \mathbf{u}_{1} & \dots & \mathbf{u}_{\rho} \\ \downarrow & & \downarrow \end{bmatrix} \begin{bmatrix} -\mathbf{s}_{1}\mathbf{v}_{1} & -\mathbf{v}_{\rho} \\ \vdots \\ -\mathbf{s}_{\rho}\mathbf{v}_{\rho} & -\mathbf{v}_{\rho} \end{bmatrix} \\
&= \begin{bmatrix} \sum_{i=1}^{\rho} u_{1i}s_{i}v_{1i} & \dots & \sum_{i=1}^{\rho} u_{1i}s_{i}v_{Li} \\ \vdots & \ddots & \vdots \\ \sum_{j=1}^{\rho} u_{Ai}s_{j}v_{1i} & \dots & \sum_{i=1}^{\rho} u_{Ai}s_{i}v_{Li} \end{bmatrix} \\
&= \begin{bmatrix} \sum_{i=1}^{\rho} s_{i}v_{1i}\mathbf{u}_{i} & \dots & \sum_{i=1}^{\rho} s_{i}v_{Li}\mathbf{u}_{i} \\ \vdots & \ddots & \vdots \\ s_{i}v_{1i}u_{Ai} & \dots & s_{i}v_{Li}u_{1i} \end{bmatrix} \\
&= \sum_{i=1}^{\rho} \begin{bmatrix} s_{i}v_{1i}u_{1i} & \dots & s_{i}v_{Li}u_{1i} \\ \vdots & \ddots & \vdots \\ s_{i}v_{1i}u_{Ai} & \dots & s_{i}v_{Li}u_{Ai} \end{bmatrix} \\
&= \sum_{i=1}^{\rho} s_{i} \begin{bmatrix} u_{1i} \\ \vdots \\ u_{Ai} \end{bmatrix} [v_{1i} \dots v_{Li}] \\
&= \sum_{i=1}^{\rho} s_{i}\mathbf{u}_{i}\mathbf{v}_{i}^{\mathsf{T}} & (A.4)
\end{array}$$

From Equation A.2 we have

$$\mathbf{x}_{\ell} = \sum_{i=1}^{\rho} s_i v_{\ell i} \mathbf{u}_i . \tag{A.5}$$

### Appendix B SVD Component Values

**Table B.1:** SVD Component  $(s_{zi}\mathbf{u}_{zi})$  Values.

		Fema	ale			Mal	е	
Age	<i>c</i> <sub>1</sub>	$c_2$	<i>c</i> <sub>3</sub>	<i>c</i> <sub>4</sub>	<i>c</i> <sub>1</sub>	$c_2$	<i>c</i> <sub>3</sub>	<i>C</i> <sub>4</sub>
0	-934.81	-32.99	-11.70	13.71	-918.91	-46.76	-6.09	5.19
1	-1078.10	-69.49	-8.07	17.13	-1067.76	-80.33	-4.90	-0.39
2	-1121.22	-67.86	-3.52	15.05	-1107.00	-74.16	-6.31	-0.64
3	-1145.29	-63.78	1.00	20.71	-1130.48	-72.86	-13.78	4.63
4	-1160.69	-58.18	-5.38	16.32	-1145.47	-65.77	-12.21	1.92
5	-1173.29	-56.67	0.27	14.96	-1156.22	-61.86	-11.92	3.09
6	-1182.52	-54.00	8.43	15.18	-1165.30	-61.35	-13.96	5.57
7	-1193.77	-53.36	14.23	13.45	-1171.97	-57.25	-14.15	5.78
8	-1201.26	-48.85	21.31	14.18	-1179.20	-53.18	-13.96	8.64
9	-1206.92	-46.74	19.06	10.84	-1185.63	-51.18	-17.16	4.86
10	-1209.42	-45.76	23.70	5.47	-1187.85	-48.52	-13.28	2.09
11	-1210.30	-46.03	19.92	-1.09	-1188.72	-44.67	-14.14	1.75
12	-1206.66	-42.32	15.42	-0.46	-1185.35	-40.50	-12.04	0.88
13	-1200.03	-40.45	13.90	-3.08	-1179.37	-38.07	-10.71	1.52
14	-1189.92	-35.59	8.62	-8.15	-1166.76	-30.08	-4.84	-3.07
15	-1178.67	-30.44	8.44	-15.37	-1153.07	-25.84	-3.74	-4.22
16	-1167.27	-25.78	0.81	-13.25	-1135.09	-17.32	-0.04	-6.87
17	-1160.91	-24.88	0.42	-10.98	-1119.99	-12.58	3.04	-6.59
18	-1154.00	-21.48	-0.52	-14.29	-1104.98	-6.06	6.08	-11.08
19	-1151.15	-22.07	-2.14	-14.19	-1098.94	-7.95	8.20	-11.61
20	-1149.00	-23.57	0.90	-12.89	-1094.97	-10.17	8.09	-12.58
21	-1148.10	-24.82	1.86	-14.46	-1092.68	-11.03	10.25	-14.70
22	-1146.33	-26.40	-0.37	-13.54	-1091.19	-12.05	11.11	-12.03
23	-1145.44	-27.81	-0.07	-14.51	-1090.64	-12.26	12.27	-12.51
24	-1143.61	-27.74	-1.07	-13.79	-1091.05	-12.59	12.21	-11.75
25	-1140.94	-25.79	-0.66	-12.45	-1090.81	-11.79	13.36	-10.84
26	-1138.44	-24.48	-3.26	-11.00	-1091.12	-11.37	13.51	-11.72
27	-1136.12	-23.62	-3.56	-8.98	-1089.91	-8.96	14.38	-10.37
28	-1134.00	-24.71	-3.27	-12.59	-1088.84	-8.40	15.09	-8.64
29	-1129.11	-19.00	-6.37	-5.78	-1087.57	-7.36	14.41	-8.24
30	-1125.27	-18.27	-5.01	-5.50	-1086.16	-9.74	13.95	-7.35
31	-1124.12	-17.81	-3.32	-7.85	-1084.21	-5.38	14.84	-7.04
32	-1119.21	-16.93	-4.52	-8.23	-1081.09	-6.38	14.14	-4.74
33	-1115.84	-15.82	-2.71	-10.07	-1078.44	-4.96	14.47	-3.97
34	-1110.86	-12.59	-7.27	-7.93	-1075.32	-3.98	14.89	-2.78
35	-1105.62	-9.14	-7.23	-5.25	-1071.97	-4.60	14.14	-1.20
36	-1102.11	-8.59	-6.12	-5.62	-1068.51	-2.62	13.42	-1.12
37	-1098.49	-7.29	-5.22	-5.60	-1065.56	-3.09	12.88	-1.85

continued on next page

Table B.1 – continued from previous page

		Fema			om previou	Male	<del></del>	
Age		<i>c</i> <sub>2</sub>	<i>c</i> <sub>3</sub>	<i>c</i> <sub>4</sub>	<i>c</i> <sub>1</sub>	<i>c</i> <sub>2</sub>	<i>c</i> <sub>3</sub>	<i>c</i> <sub>4</sub>
38	-1091.85	-2.88	-8.73	-1.87	-1060.08	-0.51	13.17	0.71
39	-1088.60	-2.96	-7.51	-3.85	-1055.88	0.89	12.19	1.07
40	-1083.01	-0.75	-7.27	1.59	-1050.70	0.45	12.45	3.47
41	-1080.12	1.69	-5.54	-4.87	-1047.16	3.79	11.64	2.54
42	-1073.06	4.63	-7.81	-0.87	-1040.78	4.13	10.94	4.04
43	-1069.13	7.38	-7.22	-0.78	-1036.22	6.42	10.83	4.10
44	-1064.54	9.28	-6.80	-2.86	-1031.37	7.15	10.00	4.58
45	-1058.85	11.16	-8.63	-0.77	-1025.18	8.52	9.46	5.31
46	-1054.89	12.11	-7.47	-2.39	-1020.74	10.15	8.92	5.73
47	-1049.45	15.59	-7.58	0.18	-1015.33	11.62	8.49	6.33
48	-1043.65	16.33	-8.00	0.91	-1009.80	11.83	7.90	6.89
49	-1038.41	18.50	-7.64	0.66	-1004.19	13.38	7.27	7.00
50	-1031.91	17.72	-9.12	0.30	-997.78	13.44	6.98	7.70
51	-1029.04	19.21	-6.45	-0.48	-994.10	15.26	6.08	7.07
52	-1021.97	18.84	-8.25	1.55	-986.90	15.74	5.85	7.90
53	-1017.78	19.66	-7.25	0.51	-982.05	16.29	4.90	7.79
54	-1012.15	20.24	-7.83	1.36	-976.10	17.54	4.57	8.11
55	-1007.32	21.09	-7.39	1.79	-970.90	18.68	4.26	7.86
56	-1001.53	21.65	-8.32	2.01	-965.39	19.28	3.29	8.29
57	-996.65	22.45	-7.52	2.27	-960.18	20.34	2.72	8.13
58	-990.38	21.82	-7.51	2.48	-953.98	20.28	2.29	8.95
59	-985.03	21.75	-6.75	2.13	-948.38	20.65	2.08	8.39
60	-977.26	20.04	-8.08	3.46	-941.20	20.05	2.62	9.34
61	-973.21	22.11	-6.90	3.08	-937.59	20.90	1.51	7.96
62	-965.41	20.87	-8.01	3.94	-930.20	20.49	1.42	8.82
63	-959.82	20.51	-5.72	3.81	-924.57	20.53	0.73	8.59
64	-953.26	20.54	-6.67	4.44	-918.57	20.66	0.63	8.62
65	-946.29	20.51	-7.38	4.65	-912.36	20.62	0.21	8.85
66	-941.19	20.99	-5.36	4.70	-907.91	21.14	-0.41	7.35
67	-934.31	21.50	-5.35	4.80	-901.75	21.18	-0.80	7.66
68	-927.11	20.83	-5.72	5.38	-895.49	21.11	-1.11	7.56
69	-920.77	21.36	-4.20	4.93	-889.83	21.20	-1.83	6.68
70	-911.67	19.74	-6.18	6.17	-882.49	19.84	-1.61	7.30
71	-907.16	21.18	-3.00	5.37	-878.42	21.09	-3.29	5.42
72	-897.89	20.17	-4.23	6.34	-870.29	20.03	-2.62	6.54
73	-891.03	20.73	-3.33	6.28	-864.21	20.13	-2.88	6.13
74	-883.26	20.99	-4.05	6.47	-857.63	19.85	-3.05	6.20
75 70	-875.74	20.88	-3.34	6.68	-851.23	19.77	-3.04	6.09
76	-868.74	21.70	-2.70	6.75	-845.16	20.05	-3.92	5.59
77	-862.35	22.39	0.07	6.19	-839.72	20.28	-5.17	3.90
78	-853.67	22.30	-2.10	6.57	-832.18	19.94	-4.29	4.94

continued on next page

Table B.1 – continued from previous page

	Female				Mal	е		
Age	<i>c</i> <sub>1</sub>	<i>c</i> <sub>2</sub>	<i>c</i> <sub>3</sub>	<i>C</i> <sub>4</sub>	<i>c</i> <sub>1</sub>	<i>c</i> <sub>2</sub>	<i>c</i> <sub>3</sub>	<i>C</i> <sub>4</sub>
79	-846.96	23.24	-0.51	5.73	-826.21	20.24	-4.63	3.72
80	-838.55	23.68	-1.82	6.12	-818.90	20.23	-3.96	4.06
81	-833.42	25.79	-0.10	4.92	-814.11	21.44	-4.44	2.57
82	-824.39	25.27	-0.03	5.26	-806.11	20.67	-4.99	2.48
83	-817.13	25.99	0.95	4.84	-799.71	20.88	-5.51	1.68
84	-809.21	26.08	1.23	4.79	-792.48	20.56	-6.08	1.48
85	-802.53	27.02	2.12	4.27	-786.78	21.46	-6.47	0.50
86	-795.34	28.10	2.95	4.05	-780.18	21.94	-7.26	-0.04
87	-788.90	29.08	3.93	3.33	-774.15	22.62	-7.74	-0.92
88	-782.19	29.71	4.56	2.95	-768.11	22.93	-8.11	-1.57
89	-775.80	31.07	5.70	2.33	-762.32	23.80	-8.66	-2.58
90	-768.10	30.58	5.38	2.39	-755.64	23.70	-8.75	-2.74
91	-763.49	32.81	7.18	1.19	-751.11	25.46	-9.64	-4.28
92	-756.27	32.87	7.39	1.19	-744.57	25.66	-10.07	-4.39
93	-750.27	33.73	8.10	0.78	-739.12	26.41	-10.58	-5.03
94	-744.09	34.24	8.51	0.53	-733.67	26.78	-11.03	-5.39
95	-737.30	34.42	9.09	0.30	-727.78	26.07	-11.10	-6.08
96	-731.39	35.07	9.66	-0.05	-722.54	26.48	-11.45	-6.59
97	-725.63	35.71	10.20	-0.40	-717.45	26.90	-11.76	-7.08
98	-720.04	36.33	10.72	-0.73	-712.52	27.30	-12.06	-7.53
99	-714.62	36.92	11.20	-1.06	-707.75	27.70	-12.32	-7.95
100	-709.39	37.49	11.65	-1.37	-703.16	28.09	-12.57	-8.34
101	-704.35	38.03	12.07	-1.66	-698.74	28.46	-12.78	-8.69
102	-699.52	38.53	12.45	-1.94	-694.50	28.81	-12.97	-9.00
103	-694.88	39.00	12.80	-2.20	-690.44	29.15	-13.13	-9.29
104	-690.46	39.43	13.11	-2.44	-686.56	29.47	-13.27	-9.53
105	-686.26	39.82	13.38	-2.67	-682.88	29.77	-13.38	-9.75
106	-682.27	40.17	13.61	-2.87	-679.38	30.05	-13.46	-9.93
107	-678.50	40.48	13.81	-3.05	-676.06	30.31	-13.53	-10.07
108	-674.95	40.74	13.97	-3.21	-672.93	30.54	-13.57	-10.19
109	-671.61	40.96	14.10	-3.36	-669.98	30.75	-13.59	-10.28

## Appendix C SVD Sum of Squares and SVD-Comp Calibration using HMD Data

The total sum of squares explained by each component of the SVD should not be interpreted like the variances identified by the eigenvalues of a PCA. Standard PCA operates on an appropriately rescaled and centered data cloud so that each new component has the same standardized scale and the 'eigenvalue' variances are real variances. In this application the SVD operates on the raw data cloud of logit-transformed mortality rates, all of which are negative. Consequently the data cloud is not centered and the dimensions of the cloud do not have the same scale. The first SVD component effectively locates the cloud with respect the origin, and because of this geometric reality, it must explain the vast majority of the total sum of squares associated with the data cloud. Another consequence is that the first component relates to the overall level of mortality and reflects the characteristic, underlying age pattern of mortality, while the remaining components describe age-specific deviations around the underlying age pattern. To better describe how much of this age-specific variability is associated with each additional component after the first, we calculate the fraction of the remaining total sum of squares associated with each of them. The 'remaining total sum of squares' is the total sum of squares remaining after subtracting the sum of squares associated with the first component. For female components 2-4, those values are 0.559472, 0.042264, and 0.034873, totaling 0.636608; and for males 0.586963, 0.073421, and 0.037134, totaling 0.697518. This indicates that the next three components after the first capture the bulk of the age-specific variation in the HMD mortality schedules. The next few components after these add little explanatory power so that it does not seem prudent to include more than four total components.

With respect to the SVs, the sum of the squares of the SVs is the total sum of squares in the original dataset (or cloud), so as either the number of points in the data cloud or the number of dimensions of the cloud increases, so will the total sum of squares and the values of the SVs, especially the first few. Consequently, the scale of the SVs is dependent on the 'size' of the dataset over which the SVD is calculated, and hence the scale of the components  $s_i \mathbf{u}_i$  is also dependent on the size of the dataset. In contrast the magnitude of the LSVs is constrained to be unity, but this means that the elements of the LSVs will be smaller as the number of elements increases, or as the number of points in the original dataset increases. All this is to explain that the scale of the components is not fixed and depends on the size of the dataset over which the SVD is calculated. Critically, this affects only the magnitude of the components, not their age patterns, and in practice none of this matters at all because the weights in Equation 11 can incorporate a factor that accounts for scale.

### **Appendix D** Estimated Regression Coefficients

**Table D.1:** Female RSV Models:  $v_{\ell i} = f_i({}_5\mathbf{q}_0\,_\ell,\,{}_{45}\mathbf{q}_{15}\,_\ell)$ 

	R	ight Singular Ve	ctor Elements	;
	<b>v</b> <sub>1</sub>	<b>v</b> <sub>2</sub>	<b>v</b> <sub>3</sub>	<b>v</b> <sub>4</sub>
<sub>5</sub> q <sub>0</sub>	0.016***	0.501***	-0.797***	1.935***
	(0.001)	(0.043)	(0.094)	(0.094)
$logit(_5q_0)$	-0.005***	-0.157***	0.202***	-0.537***
	(0.0004)	(0.013)	(0.028)	(0.028)
$logit(_5q_0)^2$	-0.001***	-0.029***	0.023***	-0.106***
	(0.0001)	(0.003)	(0.006)	(0.006)
$logit(_5q_0)^3$	-0.0001***	-0.002***	0.002***	-0.007***
	(0.00001)	(0.0002)	(0.0004)	(0.0004)
<sub>45</sub> q <sub>15</sub>	-0.003***	-0.002	0.079***	-0.049***
	(0.0001)	(0.005)	(0.010)	(0.010)
$logit(_{45}q_{15})^2$	0.0004***	0.013***	-0.024***	0.014***
	(0.00002)	(0.001)	(0.002)	(0.002)
$logit(_{45}q_{15})^3$	-0.00002***	0.002***	0.003***	0.002***
	(0.00000)	(0.0002)	(0.0004)	(0.0004)
$_{5}q_{0} \times {}_{45}q_{15}$	-0.0004***	-0.006***	0.043***	-0.004*
	(0.00002)	(0.001)	(0.002)	(0.002)
Constant	0.006***	-0.285***	0.346***	-0.934***
	(0.001)	(0.022)	(0.048)	(0.048)
Observations R <sup>2</sup> Adjusted R <sup>2</sup> F Statistic (df = 8; 4601)	4,610	4,610	4,610	4,610
	0.966	0.860	0.319	0.326
	0.966	0.860	0.318	0.324
	16 344 300***	3,542.448***	269.396***	277.570***

Note:

\*p<0.1; \*\*p<0.05; \*\*\*p<0.01

**Table D.2:** Male RSV Models:  $v_{\ell i} = f_i({}_5q_0_{\ell}, {}_{45}q_{15\ell})$ 

	Right Singular Vector Elements					
	<b>v</b> <sub>1</sub>	<b>v</b> <sub>2</sub>	<b>v</b> <sub>3</sub>	<b>v</b> <sub>4</sub>		
<sub>5</sub> q <sub>0</sub>	0.010***	0.320***	-0.425***	1.880***		
	(0.001)	(0.040)	(0.076)	(0.095)		
logit( <sub>5</sub> q <sub>0</sub> )	-0.003***	-0.109***	0.116***	-0.527***		
	(0.0003)	(0.012)	(0.023)	(0.028)		
$\log it(_5q_0)^2$	-0.001***	-0.021***	0.026***	-0.100***		
	(0.0001)	(0.002)	(0.005)	(0.006)		
$logit(_5q_0)^3$	-0.00005***	-0.002***	0.002***	-0.006***		
	(0.00000)	(0.0002)	(0.0003)	(0.0004)		
<sub>45</sub> q <sub>15</sub>	-0.002***	-0.008***	0.110***	-0.057***		
	(0.0001)	(0.003)	(0.005)	(0.007)		
$\log \mathrm{it}(_{45}\mathrm{q}_{15})^2$	0.0001***	0.002***	-0.002***	-0.004***		
	(0.00001)	(0.0004)	(0.001)	(0.001)		
logit( <sub>45</sub> q <sub>15</sub> ) <sup>3</sup>	-0.00001***	0.001***	-0.001***	-0.00004		
	(0.00000)	(0.0001)	(0.0002)	(0.0003)		
$_{5}q_{0} \times {}_{45}q_{15}$	-0.00003***	-0.001*	-0.004***	-0.002***		
	(0.00001)	(0.0004)	(0.001)	(0.001)		
Constant	0.009***	-0.194***	0.161***	-0.911***		
	(0.0004)	(0.020)	(0.039)	(0.049)		
Observations R <sup>2</sup> Adjusted R <sup>2</sup>	4,610	4,610	4,610	4,610		
	0.974	0.877	0.555	0.296		
	0.974	0.877	0.554	0.295		
Adjusted R <sup>2</sup> F Statistic (df = 8; 4601		4,095.245***	716.499***	0.295 241.669***		

Note:

\*p<0.1; \*\*p<0.05; \*\*\*p<0.01

**Table D.3:** Adult Mortality Models:  $logit(_{45}q_{15})_{z\ell} = f(_{5}q_{0\ z\ell})$ 


	logit( <sub>45</sub>	q <sub>15</sub> )
	Female	Male
<sub>5</sub> q <sub>0</sub>	-10.797*** (1.735)	2.883 (2.473)
logit( <sub>5</sub> q <sub>0</sub> )	4.005*** (0.514)	0.359 (0.740)
$logit(_5q_0)^2$	0.695*** (0.104)	0.104 (0.151)
$logit(_{5}q_{0})^{3}$	0.046*** (0.007)	0.016 (0.011)
Constant	5.921*** (0.881)	-0.703 (1.262)
Observations	4,610	4,610
$R^2$	0.934	0.793
Adjusted R <sup>2</sup>	0.934	0.793
F Statistic (df = 4; 4605)	16,192.720***	4,416.976***
Note:	*p<0.1; **p<	0.05; ***p<0.01

**Table D.4:** Infant Mortality Models:  $logit(_1q_0)_{z\ell} = f(_5q_0)_{z\ell}$ 


	logit( <sub>1</sub> q <sub>0</sub> )				
	Female	Male			
logit( <sub>5</sub> q <sub>0</sub> )	0.659*** (0.004)	0.689*** (0.004)			
$logit(_5q_0)$ $logit(_5q_0)^2$	-0.038*** (0.001)	-0.037*** (0.001)			
Constant	-0.951*** (0.006)	-0.828*** (0.006)			
Observations	4,610	4,610			
$R^2$	0.995	0.996			
Adjusted R <sup>2</sup>	0.995	0.996			
F Statistic (df = 2; 4607)	509,021.200***	569,584.700***			
Note:	*p<0.1: **	p<0.05: ***p<0.01			

*Note:* \*p<0.1; \*\*p<0.05; \*\*\*p<0.01


### Appendix E Additional Figures



**Figure E.1: Right Singular Vector Element Values for Females.** Values and predictions from model in Equation 12 on the logit scale by  $logit(_5q_0)$ . The predicted values are based on both  $_5q_0$  and  $_{45}q_{15}$  which explains why they appear as a cloud rather than a curve.



**Figure E.2: Right Singular Vector Element Values for Males.** Values and predictions from model in Equation 12 on the logit scale by  $\log \operatorname{it}(_5q_0)$ . The predicted values are based on both  $_5q_0$  and  $_{45}q_{15}$  which explains why they appear as a cloud rather than a curve.



**Figure E.3: Adult vs. Child Mortality.** Values and predictions from model in Equation 13 on the logit scale by  $logit(_5q_0)$ .






Figure E.4: Age 0 Probability of Dying  $_1\mathbf{q}_0$  vs. Child Mortality. Values and predictions from model in Equation 14 on the logit scale by  $\log \operatorname{id}_5\mathbf{q}_0$ ).



**Figure E.5: Predicted**  $_1\mathbf{q}_x$  **at Three Levels of**  $_5\mathbf{q}_0$ . As  $_5\mathbf{q}_0$  increases the relationship between female and male mortality changes, and female mortality generally exceeds male mortality between ages roughly 10 and 40 for high levels of  $_5\mathbf{q}_0$ . It has been verified that this reflects the real change in this relationship embodied in the HMD life tables.



**Figure E.6: Interquartile Range of Prediction Error by Sample Fraction.** 50 samples for each sample fraction. For each sample, the interquartile range is calculated across all ages and all mortality schedules in each sample category (in/out). Whiskers extend to 10% and 90% quantiles.

### **Appendix F** Additional Error Summary Tables

Table F.1: Weighted age-specific absolute errors in  $_5\widehat{\mathbf{q}}_x$ . Errors summed across all 4,610 HMD life tables 'SC' is SVD-Comp and 'LQ' is Log-Quad'.

		Female			Male		
x (years)	SC	LQ	SC-LQ	SC	LQ	SC-LQ	
0	1.2858	1.3096	-0.0238	1.5225	1.5418	-0.0193	
1-4	1.4056	1.2977	0.1079	2.0064	1.5467	0.4597	
5-9	0.7744	0.7395	0.0349	0.8795	0.8657	0.0138	
10-14	0.5095	0.4881	0.0215	0.5178	0.4864	0.0314	
15-19	0.6318	0.7042	-0.0724	0.8854	0.8426	0.0428	
20-24	0.7768	0.8568	-0.0800	1.6914	1.6198	0.0717	
25-29	0.7678	0.8477	-0.0798	1.5678	1.5046	0.0632	
30-34	0.7598	0.8014	-0.0416	1.5026	1.4578	0.0448	
35-39	0.8351	0.8360	-0.0009	1.6748	1.6212	0.0536	
40-44	0.9494	0.9183	0.0311	1.9518	1.8957	0.0562	
45-49	1.1049	1.0888	0.0161	2.3511	2.3162	0.0349	
50-54	1.4462	1.4425	0.0038	2.8937	2.9236	-0.0299	
55-59	1.8901	1.9089	-0.0188	3.4469	3.5936	-0.1467	
60-64	2.4265	2.5094	-0.0830	4.1203	4.4400	-0.3197	
65-69	2.9755	3.0948	-0.1194	4.5378	4.9188	-0.3810	
70-74	3.8590	4.0048	-0.1457	4.6868	5.1351	-0.4483	
75-79	4.4523	4.5111	-0.0587	4.2720	4.5634	-0.2914	
80-84	4.2565	4.3640	-0.1075	3.0786	3.2829	-0.2043	
85-89	3.0322	3.0329	-0.0007	1.7008	1.7665	-0.0657	
90-94	1.4663	1.4980	-0.0316	0.6658	0.7065	-0.0407	
95-99	0.3736	0.3983	-0.0247	0.1286	0.1440	-0.0155	
100-104	0.0460	0.0517	-0.0057	0.0133	0.0156	-0.0022	
105-109	0.0028	0.0033	-0.0005	0.0008	0.0010	-0.0001	
0-109	36.0280	36.7076	-0.6797	46.0966	47.1894	-1.0928	

**Table F.2: Weighted age-specific absolute errors in**  $\widehat{\mathbf{e}}_{X}$ **.** Errors summed across all 4,610 HMD life tables. 'SC' is SVD-Comp and 'LQ' is Log-Quad'.

	Female				Male	
x (years)	SC	LQ	SC-LQ	SC	LQ	SC-LQ
0	410.81	414.77	-3.97	618.96	630.65	-11.68
1-4	450.24	459.45	-9.21	657.75	660.59	-2.83
5-9	416.42	425.78	-9.36	608.57	633.66	-25.08
10-14	395.49	404.45	-8.96	591.07	614.95	-23.88
15-19	381.53	388.95	-7.41	580.86	604.55	-23.69
20-24	361.88	366.84	-4.95	559.07	582.08	-23.01
25-29	340.15	343.37	-3.22	520.03	544.55	-24.53
30-34	320.91	323.68	-2.77	484.94	512.07	-27.14
35-39	304.27	308.44	-4.16	450.68	479.93	-29.25
40-44	287.97	293.67	-5.70	414.35	444.17	-29.82
45-49	270.05	276.24	-6.20	374.02	402.03	-28.01
50-54	248.76	254.86	-6.10	327.74	353.29	-25.56
55-59	224.04	229.41	-5.36	275.95	297.92	-21.97
60-64	195.63	200.58	-4.95	220.36	238.34	-17.98
65-69	164.89	167.84	-2.96	163.77	176.43	-12.66
70-74	130.89	132.72	-1.82	111.93	119.33	-7.39
75-79	94.22	95.27	-1.05	67.92	70.93	-3.01
80-84	59.58	60.14	-0.56	35.90	36.53	-0.63
85-89	31.00	30.92	0.09	15.88	15.79	0.09
90-94	12.40	12.36	0.05	5.52	5.46	0.06
95-99	3.21	3.20	0.01	1.15	1.15	0.00
100-104	0.46	0.46	0.00	0.14	0.14	-0.00
105-109	0.03	0.03	0.00	0.01	0.01	0.00
110+	0.01	0.00	0.00	0.00	0.00	0.00
0+	5,104.85	5,193.42	-88.57	7,086.58	7,424.54	-337.96

Table F.3: Total Absolute Errors in  $e_0$ .

Value	Female	Male
Log-Quad	6,194	8,758
SVD-Comp, C=1	11,389	12,600
SVD-Comp, C=2	6,355	8,733
SVD-Comp, C=3	6,293	8,698
SVD-Comp, C=4	6,134	8,596
SVD-Comp, C=1 - Log-Quad	5,195	3,842
SVD-Comp, C=2 - Log-Quad	161	-25
SVD-Comp, C=3 - Log-Quad	99	-60
SVD-Comp, C=4 - Log-Quad	-59	-162