

Electronic Supplementary Material

Bacteriophages Isolated in China for the Control of *Pectobacterium carotovorum* Causing Potato Soft Rot in Kenya

Peter Muturi^{1,2}, Junping Yu¹, Alice Nyambura Maina^{4,5}, Samuel Kariuki³, Francis B Mwaura⁵, Hongping Wei^{1,6}✉

1. Key Laboratory of Emerging Pathogens and Biosafety, Wuhan Institute of Virology, Chinese Academy of Sciences, Wuhan 430071, China
2. University of Chinese Academy of Sciences, Beijing 100049, China
3. Centre for Microbiology Research, Kenya Medical Research Institute, Nairobi 00202, Kenya.
4. School of Biological and Life Sciences, Technical University of Kenya, Nairobi 00200, Kenya
5. School of Biological Sciences, University of Nairobi, Nairobi 00100, Kenya
6. Sino-Africa Joint Research Centre, Nairobi 00200, Kenya

Supporting information to DOI: 10.1007/s12250-019-00091-7




Figure S1. Blue band formation after centrifugation of phage solutions in Cesium chloride gradients. Phages were first amplified to a concentration of 1×10^9 PFUs per mL.




Figure S2. Typical spot assay of single plaque (A), clear zones on *Pectobacterium carotovorum* lawn on LB agar culture medium after infection by the isolated phages (B), determining the plaque forming units by serial dilutions of phage solutions (C), Lytic activity of *Pectobacterium* phage Wc5r on resistant *Pectobacterium carotovorum* strain (D) and *Pectobacterium atrosepticum* (E).