

Supplementary Online Content

Kharbanda EO, Vazquez-Benitez G, Lipkind HS, et al. Evaluation of the association of maternal pertussis vaccination with obstetric events and birth outcomes. *JAMA*. doi:10.1001/jama.2014.14825.

eTable 1. Clinical characteristics of women with a chorioamnionitis ICD-9 code from 2 Vaccine Safety Datalink sites, 2010-2012, chart review data

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable 1. Clinical characteristics of women with a chorioamnionitis *ICD-9* code from 2 Vaccine Safety Datalink sites, 2010-2012, chart review data

	Chorioamnionitis <i>ICD-9</i> code (658.41) N=220
	No. (%)
Clinical diagnosis of chorioamnionitis	213 (96.8)
Fever ≥ 100.4 F	179 (81.4)
Uterine or fundal tenderness	6 (2.8)
Maternal tachycardia (heart rate > 100 beats per minute)	170 (77.3)
Fetal tachycardia (heart rate > 160 beats per minute)	128 (58.2)
Malodorous vaginal fluid	7 (3.2)
Pus-like vaginal fluid	7 (3.2)
In labor prior to delivery	208 (94.6)
Prolonged labor (Labor lasting >36 hours)	2 (0.9)
Epidural	209 (95)
Preterm birth (birth < 37 weeks gestation)	14 (6.4)
Group B strep positive	49 (22.3)
Received antibiotics	200 (90.9)
Placental pathology	
Chorioamnionitis	43 (19.6)
Funisitis	16 (7.3)
Normal	5 (2.3)
Not performed	159 (72.3)