

IC	IC	F	p	η_{partial}^2	t1	p1	t2	p2	t3	p3
1	8	8.68	0.000306	0.13	0.49	0.62	-0.51	0.61	-3.74	0.000428
1	10	6.89	0.001490	0.11	-0.05	0.96	-1.26	0.21	2.41	0.019132
2	9	7.16	0.001166	0.11	-1.00	0.32	-2.05	0.05	2.65	0.010487
2	16	10.85	0.000048	0.16	-0.89	0.38	-0.51	0.61	4.51	0.000032
3	9	9.42	0.000161	0.14	-0.34	0.74	-0.42	0.68	3.68	0.000518
3	10	14.54	0.000002	0.20	-2.75	0.01	-1.44	0.15	3.47	0.000980
4	5	11.86	0.000021	0.17	-0.31	0.76	-1.90	0.06	2.72	0.008697
4	12	6.42	0.002278	0.10	-0.52	0.61	-0.90	0.37	2.07	0.043152
5	14	6.62	0.001894	0.10	-2.04	0.05	-0.08	0.94	2.28	0.026209
7	12	8.51	0.000355	0.13	0.11	0.91	1.28	0.20	5.19	0.000003
10	18	12.43	0.000013	0.18	-1.92	0.06	-0.98	0.33	4.93	0.000007
11	27	8.56	0.000342	0.13	0.22	0.83	-0.21	0.84	-3.75	0.000405
12	19	8.03	0.000543	0.12	-1.26	0.21	-0.90	0.37	3.17	0.002454
13	16	7.75	0.000694	0.12	-2.01	0.05	-0.05	0.96	3.03	0.003647
14	27	6.62	0.001900	0.10	0.50	0.62	-2.13	0.04	2.26	0.027408
16	17	8.81	0.000273	0.13	0.91	0.36	0.52	0.60	-4.19	0.000097
16	22	7.03	0.001313	0.11	0.22	0.83	1.11	0.27	-2.87	0.005647

FDR level:	0.002278	No significant edges	No significant edges	0.0432
-------------------	-----------------	-----------------------------	-----------------------------	---------------

SI-Table 1: Results from the repeated measures ANOVA for the 17 edges showing a significant group x time interaction.

FDR correction was applied to each test individually.