

#### **Supplementary file 4A.**

Influence analysis of prevalence of intradialytic hypotension defined according to the EBPG definition on session level (random effects model)

	Proportion	95% confidence interval
<b>Omitting Akhmouch, 2010</b>	12.0	6.9-20.2
<b>Omitting Flythe Hemocohort, 2014</b>	10.3	4.7-21.2
<b>Omitting Kuipers, 2016</b>	11.2	6.0-20.0
<b>Omitting Rocha, 2015</b>	7.4	5.2-10.4
<b>Omitting Steinwandel, 2018</b>	10.3	0.06-0.19
<b>Pooled estimate (without omitting studies)</b>	0.10	5.5-18.6

#### **Supplementary file 4B.**

Influence analysis of prevalence of intradialytic hypotension defined according to the SBP nadir <90 mmHg definition on session level (random effects model)

	Proportion	95% confidence interval
<b>Omitting Cho, 2017</b>	0.12	0.12-0.12
<b>Omitting Flythe Hemocohort, 2014</b>	0.12	0.12-0.12
<b>Omitting Flythe/ LDO, 2014</b>	0.16	0.16-0.16
<b>Omitting Orofino, 1990</b>	0.13	0.12-0.12
<b>Omitting Sands, 2014</b>	0.10	0.10-0.10
<b>Pooled estimate (without omitting studies)</b>	0.12	0.12-0.12