

Supplementary Table 2. Baseline characteristics of patients according to the occurrence of death in 383 patients

Variable	No death (n = 358)	Death (n = 25)	p value
Age, yr	55.8 ± 13.0	65.9 ± 10.7	< 0.001
Male sex	236 (65.9)	16 (64.0)	0.830
ESRD cause			
Diabetic	178 (49.7)	17 (68.0)	0.098
BMI, kg/m ²	24.7 ± 3.7	23.9 ± 2.6	0.285
SBP, mmHg	129.2 ± 14.5	133.2 ± 16.0	0.199
DBP, mmHg	75.5 ± 10.0	78.0 ± 8.1	0.242
Serum Cr, mg/dL	2.80 ± 0.66	3.00 ± 0.79	0.163
eGFR, mL/min/1.73 m ²	27.15 ± 7.12	22.24 ± 6.60	0.001
Urinary protein, g/g Cr	2.12 ± 2.34	1.88 ± 2.11	0.617
Hemoglobin, g/dL	11.4 ± 1.7	10.4 ± 1.7	0.005
Albumin, g/dL	4.0 ± 0.4	3.9 ± 0.5	0.243
Uric acid, mg/dL	8.4 ± 1.9	8.0 ± 1.9	0.396
LDL, mg/dL	92.5 ± 35.5	91.2 ± 20.2	0.852
CRP, mg/dL	0.68 ± 2.80	0.52 ± 0.96	0.788
Calcium, mg/dL	8.87 ± 0.57	8.78 ± 0.74	0.476
Phosphorous, mg/dL	3.89 ± 0.73	3.88 ± 0.83	0.962
Renalase, µg/mL	74.7 ± 34.2	92.4 ± 40.3	0.014
Follow-up period, mon	57.8 ± 14.8	34.2 ± 17.4	0.004

Values are presented as mean ± SD or number (%).

ESRD, end stage renal disease; BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; Cr, creatinine; eGFR, estimated glomerular filtration rate; LDL, low density lipoprotein; CRP, C-reactive protein.