

Supplementary Table 3. Baseline characteristics of patients according to the occurrence of renal outcomes in 322 patients

Variable	No renal outcome (n = 185)	Renal outcome (n = 137)	p value
Age, yr	57.6 ± 13.3	54.1 ± 12.7	0.016
Male sex	121 (65.4)	90 (65.7)	1.000
ESRD cause			
Diabetic	80 (43.2)	82 (59.9)	0.003
BMI, kg/m ²	24.8 ± 3.7	24.3 ± 3.4	0.242
SBP, mmHg	128.5 ± 15.0	130.6 ± 14.0	0.212
DBP, mmHg	75.1 ± 9.6	76.2 ± 9.9	0.292
Serum Cr, mg/dL	2.60 ± 0.55	3.03 ± 0.69	< 0.001
eGFR, mL/min/1.73 m ²	28.29 ± 7.26	25.21 ± 5.95	< 0.001
Urinary protein, g/g Cr	1.27 ± 1.54	3.18 ± 2.21	< 0.001
Hemoglobin, g/dL	11.8 ± 1.8	10.9 ± 1.5	< 0.001
Albumin, g/dL	4.1 ± 0.4	3.8 ± 0.5	< 0.001
Uric acid, mg/dL	8.3 ± 1.8	8.3 ± 1.8	0.749
LDL, mg/dL	91.6 ± 26.0	93.1 ± 33.3	0.677
CRP, mg/dL	0.69 ± 2.65	0.61 ± 3.09	0.823
Calcium, mg/dL	9.00 ± 0.55	8.72 ± 0.59	< 0.001
Phosphorous, mg/dL	3.75 ± 0.70	4.10 ± 0.69	< 0.001
Renalase, µg/mL	69.6 ± 31.8	84.0 ± 35.8	< 0.001
Follow-up period, mon	32.4 ± 4.2	21.5 ± 8.5	< 0.001

Values are presented as mean ± SD or number (%).

ESRD, end stage renal disease; BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; Cr, creatinine; eGFR, estimated glomerular filtration rate; LDL, low density lipoprotein; CRP, C-reactive protein.