

Supplemental Table 1. Multivariable predictors of LV concentric remodelling (LV mass/volume)

	Standardised coefficient (β)	95% CI	p-value
Age	0.136	0.000 – 0.005	0.091
Gender	0.367	0.069 – 0.165	<0.001
Ethnicity	0.240	0.035 – 0.161	0.003
T2D	0.194	0.009 – 0.115	0.023
Hypertension	0.139	-0.006 – 0.159	0.070
BMI	0.153	0.000 - 0.007	0.070

Abbreviations: T2D=type 2 diabetes, BMI=body mass index.

Supplemental Table 2. Correlations between plasma biomarkers, LV mass/volume and LA volume.

	LV mass/volume		Maximal indexed LA volume	
	Pearson's correlation coefficient (r)	p-value	Pearson's correlation coefficient (r)	p-value
FABP4	0.082	0.366	-0.146	0.096
Adiponectin	-0.328	<0.001	0.291	0.001
Interleukin-6	0.063	0.486	0.034	0.700
Interleukin-8	0.024	0.791	0.043	0.623
TNFR1	0.249	0.005	-0.018	0.841
SerpinE1 PAI1	0.073	0.416	-0.352	<0.001
GDF15	0.273	0.002	0.052	0.554
MMP7	0.221	0.013	0.002	0.983
TIMP1	0.079	0.381	-0.066	0.456

Abbreviations: FABP4=fatty acid binding protein 4, TNFR1=tumour necrosis factor receptor 1, Serpin E1 PAI1=SerpinE1 plasminogen activator inhibitor 1, GDF15=growth differentiation factor 15, MMP7=matrix metalloproteinase 7, TIMP1=tissue inhibitor of metalloproteinases 1.