

# A mathematical model to predict mean time to delivery following cervical ripening with dinoprostone vaginal insert.

**Fanny Levast<sup>1, \*</sup>, Guillaume Legendre<sup>1</sup>, Hady El Hachem<sup>2</sup>, Patrick Saulnier<sup>3</sup>, Philippe Descamps<sup>1</sup>, Philippe Gillard<sup>1</sup>, Pierre-Emmanuel Bouet<sup>1</sup>**

<sup>1</sup> Department of Obstetrics and Gynecology, Angers University Hospital, Angers, France

<sup>2</sup> Department of Reproductive Medicine, Clemenceau Medical Center, Beirut, Lebanon

<sup>3</sup> Department of Methodology and Biostatistics, Angers University Hospital, Angers, France

\*Correspondence and requests for materials should be addressed to F.L (email: [fannylevast@yahoo.fr](mailto:fannylevast@yahoo.fr))

**Table of statistics of residuals (a)**

	Minimum	Maximum	Mean	Standard deviation	N
Predictors	710,5430	1893,6970	1236,7085	216,57145	398
Residuals	-1295,73462	2382,65210	,00000	616,30556	398
Standardized predictors	-2,430	3,034	,000	1,000	398
Standardized residuals	-2,092	3,846	,000	,995	398

"a" Dependent variable: Time to delivery