

Supporting Information

Recovery of polyphenols from grape pomace using polyethylene glycol (PEG)-grafted silica particles and PEG assisted co-solvent elution

*Ayca Seker ^{*1}, Baran Arslan ², and Shulin Chen ¹*

¹Department of Biological Systems Engineering, Washington State University, Pullman, WA, 99164-6120, United States

²Gene and Linda Voiland School of Chemical Engineering and Bioengineering, Washington State University, Pullman, WA, 99164-6515, United States

Table S1. Individual phenolic compounds of grape pomace extract

	Individual Polyphenols (mg L ⁻¹)									
	GA ¹	CA ²	PB2 ³	DC ⁴	pCA ⁵	IC ⁶	MC ⁷	MY ⁸	QU ⁹	KA ¹⁰
Grape pomace extract	70.2 ±0.6	983.2 ±37.7	288.8 ±2.5	486.7 ± 31.2	47.9 ±5.4	487.3 ±0.9	345.6 ±3.1	10.4± 0.5	275.3 ±0.4	259.4 ±4.7

¹GA:Gallic Acid, ²CA:(+)-Catechin, ³PB2:Procyanidin-B2, ⁴DC:Delphinidin Chloride, ⁵pCA:P-coumaric acid, ⁶IC:Isoquercetin, ⁷MC: Malvidin Chloride, ⁸Myricetin: MY, ⁹Quercetin: QU, ¹⁰Kaempferol: KA. Data are represented in mean ± standard deviation of duplicates.