

Supplementary Online Content

Alore EA, Suliburk JW, Ramsey DJ, et al. Diagnosis and management of primary hyperparathyroidism across the Veterans Affairs health care system. *JAMA Intern Med*. Published online July 15, 2019.

doi:10.1001/jamainternmed.2019.1747

eTable. Comorbid Conditions Included in the Elixhauser Comorbidity Index


eFigure. Indications for Parathyroidectomy by Consensus Guidelines Over Time

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Comorbid Conditions Included in the Elixhauser Comorbidity Index

1	AIDS	16	Liver disease
2	Alcohol abuse	17	Lymphoma
3	Blood loss anemia	18	Metastatic cancer
4	Cardiac arrhythmias	19	Obesity
5	Chronic pulmonary disease	20	Other neurological disorders
6	Coagulopathy	21	Paralysis
7	Congestive heart failure	22	Peptic ulcer disease excluding bleeding
8	Deficiency anemias	23	Peripheral vascular disorders
9	Depression	24	Psychoses
10	Diabetes, complicated	25	Pulmonary circulation disorders
11	Diabetes, uncomplicated	26	Renal failure
12	Drug abuse	27	Rheumatoid arthritis/ collagen vascular diseases
13	Fluid and electrolyte disorders	28	Solid tumor without metastasis
14	Hypertension	29	Valvular disease
15	Hypothyroidism	30	Weight loss

eFigure. Indications for Parathyroidectomy by Consensus Guidelines Over Time



Abbreviations: NIH = National Institutes of Health; DEXA = Dual-energy X-ray absorptiometry; pHPT = primary hyperparathyroidism