

| <i>Orthoptic examination parameters</i> | M3 vs M0 | | M9 vs M0 | | M9 vs M3 | |
|----------------------------------------------------|------------------------------|-------------------|------------------------------|-------------------|------------------------------|----------------|
| NPC (cm) | -3.26 [-4.15 ; -2.37] | p<.0001 | -3.23 [-4.16 ; -2.30] | p<.0001 | 0.03 [-0.57 ; 0.63] | p=0.92 |
| far convergence amplitude (prism diopters) | 20.04 [15.82 ; 24.26] | p<.0001 | 15.18 [11.59 ; 18.77] | p<.0001 | -4.86 [-7.96 ; -1.76] | p=0.004 |
| Near convergence amplitude (prism diopters) | 14.54 [10.39 ; 18.68] | p<.0001 | 11.19 [7.23 ; 15.15] | p<.0001 | -3.34 [-5.11 ; -1.57] | p=0.001 |
| far divergence amplitude (prism diopters) | 0.64 [0.01 ; 1.26] | p=0.06 | 0.62 [-0.02 ; 1.26] | p=0.07 | -0.02 [-0.48 ; 0.44] | p=0.94 |
| Near divergence amplitude (prism diopters) | 0.92 [-0.13 ; 1.97] | p=0.12 | 0.53 [-0.68 ; 1.74] | p=0.67 | -0.39 [-1.25 ; 0.46] | p=0.37 |

Table 3
ORTE