

Supporting Information

**Indolyl-Pyridinyl-Propenone-Induced Methuosis
through the Inhibition of PIKFYVE**

Hyelim Cho,^{*,†,§} Erin Geno,^{†,||} Maude Patoor,[†] Adam Reid,[‡] Richard I. McDonald,[†] Marc Hild,^{*,†} and Jeremy L. Jenkins^{*,†}

[†]Chemical Biology and Therapeutics, Novartis Institutes for BioMedical Research, 181 Massachusetts Avenue, Cambridge, Massachusetts 02139, USA

[‡]Department of Chemistry, Johns Hopkins University, 3400 N. Charles Street, Baltimore, Maryland 21218, USA

Figure S1. Synthetic Protocols






Figure S2. ^1H NMR spectra


^1H NMR (400 MHz, DMSO-d6) of **1a**




¹H NMR (400 MHz, DMSO-d6) of **2a**



¹H NMR (400 MHz, DMSO-d6) of **2b**




¹H NMR (400 MHz, DMSO d6) of **2c**



S5

¹H NMR (400 MHz, DMSO-d6) of **2l**



¹H NMR (400 MHz, DMSO-d6) of **2o**






Figure S3. HPLC spectra


HPLC of **1a**




HPLC of **2a**




HPLC of **2b**



HPLC of 2c



HPLC of 2l



HPLC of 2o







Figure S4. HR-MS spectra




HR-MS of 2a




HR-MS of 2b



HRMS of 2c



HR-MS of **2l**



HR-MS of **2o**

