

## ONLINE RESOURCE 2: SUPPLEMENTARY TABLE 1.

Supplementary Table 1. Parameter estimates and average posterior probabilities for the three trajectories of postpartum depressive symptoms.

Trajectory of postpartum depressive symptoms	N (%)	APP	Parameters	B (SE)
1. Low symptoms trajectory	1309 (91.7)	.97	<i>Mean</i>	
			Intercept	4.08 (0.17)***
			Linear	-0.03 (0.006)***
			Quadratic	0.00 (0.00)**
			<i>Variance</i>	
			Intercept	11.00 (1.09)***
2. Increasing-decreasing symptoms trajectory	77 (5.4)	.80	<i>Mean</i>	
			Intercept	11.93 (0.73)***
			Linear	0.38 (0.08)***
			Quadratic	-0.01 (0.002)***
			<i>Variance</i>	
			Intercept	11.00 (1.09)***
3. Increasing symptoms trajectory	41 (2.9)	.83	<i>Mean</i>	
			Intercept	7.07 (0.99)***
			Linear	0.20 (0.09)*
			Quadratic	0.00 (0.002)
			<i>Variance</i>	
			Intercept	11.00 (1.09)***

APP, average posterior probability; SE, standard error. For the intercept growth factor, the variance was estimated but held equal across classes. For the linear (slope) and quadratic growth factors, the variance was fixed at zero in each class. \* $p < .05$ , \*\* $p < .01$ , \*\*\* $p < .001$ .

## Obsessive-compulsive personality disorder symptoms as a risk factor for postpartum depressive symptoms

*Archives of Women's Mental Health*

Kiki E.M. van Broekhoven, Annemiek Karreman, Esther E. Hartman, Paul Lodder, Joyce J.

Endendijk, Veerle Bergink, Victor J.M. Pop<sup>1</sup>. <sup>1</sup>Corresponding author. Email address:

[v.j.m.pop@uvt.nl](mailto:v.j.m.pop@uvt.nl). Department of Medical and Clinical Psychology, Tilburg University.