## **Supporting Information**

## Decarbonylative Borylation of Amides by Palladium Catalysis

Shicheng Shi,† and Michal Szostak\*,†


<sup>†</sup>Department of Chemistry, Rutgers University, 73 Warren St., Newark, NJ 07102, United States

michal.szostak@rutgers.edu

Table of Contents	<b>S</b> 1
<sup>1</sup> H and <sup>13</sup> C NMR Spectra	S2

## **Corresponding Author:**

Prof. Dr. M. Szostak Department of Chemistry, Rutgers University 73 Warren Street, Newark, NJ 07102, United States E-mail: michal.szostak@rutgers.edu







Figure S1.  $^{1}$ H NMR and  $^{13}$ C NMR Spectra of 2a.







Figure S2.  $^{1}$ H NMR and  $^{13}$ C NMR Spectra of 2b.









Figure S3.  $^{1}$ H NMR and  $^{13}$ C NMR Spectra of 2c.





**Figure S4.** <sup>19</sup>F NMR Spectrum of **2c**.







Figure S5. <sup>1</sup>H NMR and <sup>13</sup>C NMR Spectra of 2d.









Figure S6. <sup>1</sup>H NMR and <sup>13</sup>C NMR Spectra of 2e.







Figure S7. <sup>1</sup>H NMR and <sup>13</sup>C NMR Spectra of 2f.







Figure S8. <sup>1</sup>H NMR and <sup>13</sup>C NMR Spectra of 2g.






Figure S9. <sup>1</sup>H NMR and <sup>13</sup>C NMR Spectra of 2h.




Figure S10. <sup>1</sup>H NMR and <sup>13</sup>C NMR Spectra of 2i.




Figure S11. <sup>1</sup>H NMR and <sup>13</sup>C NMR Spectra of 2j.