

Review

Significance of Vectorcardiogram in the Cardiological Diagnosis of the 21st Century

ANDRÉS RICARDO PÉREZ RIERA, M.D., AUGUSTO H. UCHIDA, M.D.,* CELSO FERREIRA FILHO, M.D.,[†]
ADRIANO MENEGHINI, M.D.,[‡] CELSO FERREIRA, M.D., PH.D.,[¶] EDGARDO SCHAPACKNIK, M.D.,[§] SERGIO DUBNER, M.D.,**
PAULO MOFFA, M.D., PH.D.^{††}

Chief of the Sector of Electrovectorcardiography of the Discipline of Cardiology, School of Medicine, ABC Foundation, Santo André; *Assistant MD of the Electrocardiology Division, Heart Institute, University of Sao Paulo Medical School, [†]Full Professor of the School of Medicine of Santo Amaro, UNISA, Assistant Professor of the Discipline of Cardiology, School of Medicine, ABC Foundation, Santo André, [‡]Full-time Assistant Professor of the Discipline of Cardiology, School of Medicine, ABC Foundation, Santo André, [¶]Full Professor of the Discipline of Cardiology, School of Medicine, ABC Foundation, Santo André, “Livre Docente” Professor of the Federal University of São Paulo, São Paulo, Brazil; [§]Chief of the Department of Chagas Disease of the Dr. Cosme Argerich Hospital, **Director of Arrhythmias and Electrophysiology Service, Clinical and Maternidad Suizo Argentina, Buenos Aires, Argentina; ^{††}Associate Professor of Cardiology and Director of the Electrocardiology Division, Heart Institute, University of Sao Paulo Medical School, São Paulo, Brazil

Summary

Until the mid-1980s, it was believed that the vectorcardiogram presented a greater specificity, sensitivity and accuracy in comparison to the conventional electrocardiogram, in the diagnosis of the different heart diseases. Recent studies revealed that the vectorcardiogram still is superior to the electrocardiogram in very specific situations, such as in the evaluation of electrically inactive areas, in intraventricular conduction disorders combined and/or in association to inactive areas, in the identification and location of ventricular preexcitation, in the differential diagnosis of patterns varying from normal of electrical axis deviation, in the evaluation of particular aspects of Brugada syndrome, and in the estimation of the severity of some enlargements, among others.

With the advent of computerized vectorcardiography, a technology that improves the processing and recording method; a future still promising is expected for this methodology.

In the fields of education and research, vectorcardiography provided a better and more rational insight into the electrical phenomena that occurs spatially, and represented an important impact on the progress of electrocardiography. Although a few medical centers still use the method as a routine, we hope that the use of this resource will not get lost over time, since vectorcardiography still represents a source to enrich science by enabling a better morphological interpretation of the electrical phenomena of the heart.

Key words: vectorcardiogram, electrocardiogram, cardiovascular diagnostic technic, sensitivity, specificity

Address for reprints:
Sergio Dubner, M.D.
Director of Arrhythmias and Electrophysiology Service
Clinical and Maternidad Suizo Argentina
Arenales 2463 3 A
1124 Buenos Aires
Argentina
e-mail: dubner@ciudad.com.ar

Received: December 13, 2005
Accepted with revision: February 27, 2006

Published online in Wiley InterScience
(www.interscience.wiley.com).
DOI:10.1002/clc.14
© 2007 Wiley Periodicals, Inc.

Clin. Cardiol. 2007; 30: 319–323.
© 2007 Wiley Periodicals, Inc.

Concept

The vectorcardiogram or VCG (Fig. 1) is the spatial representation of electromotive forces generated during cardiac activity and is analyzed in three spatial planes (horizontal, frontal and sagittal).¹

An instantaneous electric dipole is formed each moment during ventricular depolarization. The addition of all individual dipoles generates the resulting dipole




FIG. 1 P, QRS and T—Loops of vectorcardiogram on three planes.

of the cardiac electrical activity, moment to moment, represented by a vector.

Spatial vectorcardiography is the form of electrocardiography that tries to describe the electromotive force developed by the heart each instant as a single vector, while all the successive instantaneous vectors have a common point of origin.²

Vector

A Measurement unit that has direction or orientation and module, magnitude and intensity, used in electrovectorcardiography to represent the dipole of depolarization and repolarization. All vectors have an onset and an end called origin and end.

The size of a vector determines the magnitude, the orientation and the direction in the electric field represented by it, while the point of the vector indicates its positive side. Therefore, vectocardiographic loops represent the position of all the instantaneous vectors, at each moment, during cardiac repolarization, obtaining different loops for the P, QRS, T and U waves.¹

Vectorcardiography is based on the concept of the dipole as an approximation equivalent originating in the heart, and uses corrected orthogonal leads, which determine three spatial planes (Fig. 2): frontal plane (FP), horizontal plane (HP) and left sagittal plane (LSP) or




FIG. 2 The three orthogonal leads and the three planes on vectorcardiography.

right sagittal plane (RSP). The term orthogonal originates in the fact that the axes of the three planes are perpendicular to each other, and corrected because technical devices of resistance and multiple connections that correct the deficient homogeneity of the electric field that surrounds the heart are used. These three corrected orthogonal leads of Frank's system, as well as the three planes determined by them, cross each other at a central point called E point, thus forming a 90° angle with each other.

Conventionally, the horizontal lead that extends from left (0°) to right ($+/-180^\circ$) is called X. The axis of the corrected X lead corresponds approximately to the bipolar DI lead and the V_6 precordial lead. This lead forms the HP and the FP.

The vertical lead is known as the Y orthogonal lead, and it stretches from down ($+90^\circ$) to the top (-90°) and it approximately corresponds to the unipolar aVF lead of the electrocardiogram (ECG), which has its positive pole in $+90^\circ$. It provides information about the inferosuperior orientation of the vectors. The Y lead forms the FP and the LSP or the RSP.

Finally the axis of the sagittal orthogonal lead known as the Z axis, stretches from the back ($+90^\circ$) to the front (-90°) or posteroanterior orientation, with its posterior part being positive and its anterior part being negative. The Z orthogonal lead corresponds approximately to the precordial V_2 lead of the conventional ECG and it forms the HP and the LSP or the RSP.

Advantages of the VCG Compared to the ECG

1. The vectorcardiogram (VCG) provides three-dimensional information of the electric activity of the atria and the ventricles, showing in a clearer way than the ECG, the spatial orientation and the magnitude of the vectors at every moment.³

2. The VCG has a greater sensitivity than the ECG in detecting atrial enlargements⁴ and greater sensitivity and specificity than the ECG in the diagnosis of left ventricular enlargement (LVE). Abbott-Smith *et al.* made vectorcardiograms in 100 patients carriers of LVE confirmed in the necropsy⁵ and concluded that the VCG was capable of diagnosing 50% of the cases, with 11.7% of false positives.
3. The VCG may clear doubts in the cases of suspicion of electrically inactive area in the septal or anteroseptal wall of the left ventricle (LV), when the LVE of the systolic type is present, observed in ECG with QS pattern in V1; V1 and V2 or V1, V2 and V3. In the absence of an electrically inactive area, the “dashes” of the initial 10 to 20 ms of the QRS loop, are recorded without delay, while in the presence of electrically inactive area, the dashes of the initial 40 ms are very close to each other.⁶
4. The VCG presents a greater correlation with the echocardiogram, when compared with the ECG, in determining the left ventricular mass,⁷ and it appears to be superior to the ECG and echocardiogram in the diagnosis of chamber enlargement, associated with electrically inactive areas.⁸
5. The VCG presents a greater diagnostic sensitivity in comparison to the ECG in acute myocardial infarction (AMI), when associated with left anterior fascicular block (LAFB). In the presence of AMI of the LV inferior wall, the VCG may bring additional information, which the ECG does not reveal, such as the association with LAFB.⁹
6. The VCG presents a greater sensitivity and specificity than the ECG in the diagnosis of strict dorsal AMI, and it enables a more appropriate differentiation with other causes of prominent anterior forces, such as normal hearts with counterclockwise rotation of the longitudinal axis and shift to the right of the transition area in precordial leads, right ventricle enlargement, complete right bundle branch block (CRBBB), hypertrophic cardiomyopathy both in its obstructive and nonobstructive form (increase in the magnitude of the septal vector), diastolic enlargement of the left ventricle with dislocation of the transition area to the right, ventricular pre-excitation of the Wolff-Parkinson-White type (WPW type) with anomalous bundle, in a parallel way to posterior location WPW type A, Duchenne-Erb myopathy or malignant in childhood and other causes.^{10,11}
7. The VCG has more sensitivity than the ECG for the diagnosis of multiple infarctions, associated with LAFB.¹²
8. The VCG has a greater accuracy than the ECG in the diagnosis of inferior infarction,⁹ however, there is no consensus since there are studies that concluded that the VCG is not superior to the ECG in the diagnosis of isolated diaphragmatic infarction.¹³ Edenbrandt *et al.* compared the diagnostic value of both methods in 65 patients with inferior AMI proven by hemodynamic study and gammagraphy with Thallium 201, observing that the sensitivity of the VCG was 69%, and the ECG was 43%, with $p < 0.001$. The control group showed three false positives.¹⁴
9. The method improves sensitivity in the diagnosis of inferior infarction extended to the LV anterior wall.¹⁵
10. The VCG is of great significance for the diagnosis of the left septal fascicular block (LSFB).^{16–18} This type of left fascicular block was shown in numerous publications and the Anglo-Saxon literature still does not acknowledge it.
11. The VCG is superior to the ECG in the cases of atypical CRBBB associated with LAFB (bifascicular block) called by Rosenbaum as the “standard masquerading bundle branch block”. In these cases, in the presence of CRBBB associated to a high degree of LAFB, the DI lead presents small or nonexistent S wave, with a pure R wave appearing in this lead, characteristic of complete left bundle branch block (CLBBB) (pseudo CLBBB). This situation translates the presence of CRBBB associated to LAFB, LVE and block located in the left ventricle.¹⁹ In some cases, a CRBBB pattern is observed in the right precordial leads and CLBBB in the left precordial leads. This situation was called “masquerading bundle branch block”. This pattern defines the presence of CRBBB associated with severe LVE, a block located in the anterolateral wall of the left ventricle and usually LAFB.²⁰
12. The VCG is very useful to differentiate the rare CLBBB with extreme deviation of SAQRS to the right in the FP (to the right of $+90^\circ$). According to the location of SAQRS in the FP, the CLBBB was divided into 4 types:
 1. CLBBB with SAQRS not deviated: between -30° and $+60^\circ$. It represents 65–70% of the cases;
 2. CLBBB with SAQRS with extreme deviation to the left: beyond -30° . It represents 5% of the total;
 3. CLBBB with SAQRS deviated to the right between $+60^\circ$ and $+90^\circ$. It represents 4% of the total;
 4. CLBBB with SAQRS presenting extreme deviation to the right: $> +90^\circ$. This group represents less than 1% of the total of CLBBB and was called “paradoxical type” by Lepschkin.
 CLBBB with SAQRS located to the right of $+90^\circ$ in the FP, may have SAQRS located in

the right inferior or right superior quadrant. Lepeschkin called them "paradoxical CLBBB" or type IV (SAQRS between $+90^\circ$ and $+135^\circ$). We could add a type V when SAQRS is located to the right of $+135^\circ$ (CLBBB of congenital heart diseases). In these cases, the VCG is superior to ECG in determining the possible cause:

1. If CLBBB is associated with severe right ventricular enlargement (RVE);
 2. If fascicular CLBBB (LAFB + LPFB) by a higher degree of block in the left posterior fascicle;
 3. If the CLBBB is associated with lateral electrically inactive areas.
13. The technique known as Continuing Vectorcardiography Monitoring (CVM) carried out during elective angioplasty, proved to be a promising tool to detect patients with an increased risk of developing AMI related to the procedure. Guo *et al.*²¹ used the method in 169 patients, which started 5 min before the procedure and was interrupted 30 min after the first insufflation of the angioplasty balloon. Considering the ST segment elevation to determine the AMI, the sensitivity of the CVM to detect increased risk of acute myocardial infarction related to the procedure was 93%, the specificity was 56% and the negative predictive value 99%.
 14. The VCG presents a greater diagnostic sensitivity than ECG to determine the severity of congenital aortic valve stenosis. Thus, the presence of the maximal vector in the horizontal plane to the left maximal spatial voltage (LMSV) with a voltage greater than 4 mV, heading to the left and backward around -56° , represents a significant marker of severe aortic stenosis (left intraventricular pressure >200 mmHg); the presence of the maximal vector to the left with a voltage near 2.2 mV and around -19° , indicates mild congenital aortic stenosis.²²
 15. In patient carriers of congenital pulmonary valve stenosis, the VCG has a good correlation between the value of the systolic pressure of the right ventricle and the presence of the maximal spatial vector to the right of the HP: "Maximal Spatial Voltage directed to the Right" (RMSV). Thus, a right intraventricular pressure >100 mmHg has a RMSV >2.3 mV.²³
 16. The VCG is superior to the ECG to identify and locate the anomalous bundle in pre-excitation of the Wolff-Parkinson-White. The method presents a high sensitivity and accuracy. This fact is relevant to guide the electrophysiologist, pointing the most appropriate site to apply radiofrequency energy.²⁴ The diagnostic specificity is not increased when compared to an ECG in this case.³
 17. The VCG presents greater sensitivity and specificity than the ECG in the diagnosis of end conduction delay by the fascicles of the right branch (blocks of the right branch: fascicular, zonal or of the free wall). The VCG enables to rule out or confirm the cases where the ECG presents a doubt when there is association of end delay through the right branch with electrically inactive areas, both of the inferior and the anterior walls.²⁵
 18. The VCG optimizes the differential diagnosis of right fascicular blocks with left fascicular blocks.²⁶
 19. The vectorcardiogram is very useful in the diagnosis of Brugada syndrome when the ECG shows extreme deviation of SAQRS to the left in the FP (9.5% of the cases).²⁷ We showed that in this entity, the extreme deviation of SAQRS to the left might be the consequence of LAFB and of end conduction delay through the superior or subpulmonary fascicle of the right branch, which goes through the right ventricle outflow tract, the area affected in this entity.²⁸
 20. The VCG has a great value in the analysis of electrical modifications that are the consequence of septal percutaneous ablation of the obstructive form of severe hypertrophic cardiomyopathy, not responsive to drugs and with incapacitating symptoms (functional class II and IV). The result of septal or anteroseptal infarction generates a pattern of CRBBB in almost all cases, unlike myotomy/myectomy surgery, which promotes CLBBB in approximately 80% of the cases.²⁹

With the use of computerized VCG, obtaining and processing graphs is easier, and the problems of measuring the loops are eliminated, since it is possible to determine where each one begins and ends, establishing in a precise way, the ratio of length and width of T waves, and the estimation of the areas of the loops. In comparison with the traditional recording method, computerized VCG has a greater accuracy in measurement, besides a great processing velocity.^{21,30}

In spite of the studies that show that the VCG and the ECG have a very similar diagnostic capacity,³¹ the VCG is still evolving and it will always have didactic usefulness to teach electrocardiology, besides representing a low-cost method, with great diagnostic value in different situations where electrocardiographic recording is doubtful.^{32,33}

References

1. Grishman A, Donoso E. Spatial vectorcardiography I. *Mod Concepts Cardiovasc Dis* 1961;30:687-692

2. Helm RA. Theory of vectorcardiography: a review of fundamental concepts. *Am Heart J* 1955;49:135–159
3. Chou TC. Value and limitations of vectorcardiography in cardiac diagnosis. *Cardiovasc Clin* 1975;6:163–178
4. Chou TC. When is the vectorcardiogram superior to the scalar electrocardiogram? *J Am Coll Cardiol* 1986;8:791–799
5. Abbott-Smith CW, Chou T. Vectorcardiographic criteria for the diagnosis of left ventricular hypertrophy. *Am Heart J* 1970;79:361–369
6. Pipberger HV, Goldman MJ, Littmann D, Murphy GP, Cosma J, et al.: Correlations of the orthogonal electrocardiogram and vectorcardiogram with constitutional variables in 518 normal men. *Circulation* 1967;35:536–551
7. Bocanegra Arroyo J, Braga JMS, Luna Filho B. Análise crítica do eletrocardiograma e do vetocardiograma no diagnóstico da hipertrofia ventricular esquerda. *Rev Soc Cardiol Estado de São Paulo* 1994;4:353–360
8. Vine DL, Finchum RN, Dodge HT. Comparison of the vectorcardiogram with the electrocardiogram in the prediction of left ventricular size. *Circulation* 1971;43:547–558
9. Hurd HP II, Starling MR, Crawford MH. Comparative accuracy of electrocardiographic and vectorcardiographic criteria for inferior myocardial infarction. *Circulation* 1981;63:25–1029
10. Brisse B. Clinical vectorcardiography: the Fritz-Schellong commemorative lecture. *Z Kardiol* 1987;76:65–71
11. Hoffman I, Taymor RC, Morris MH, Kittell I. Quantitative criteria for the diagnosis of dorsal infarction using the Frank Vectorcardiogram. *Am Heart J* 1965;70:295–304
12. Benchimol A, Desser KB. Advances in clinical vectorcardiography. *Am J Cardiol* 1975;36:76–86
13. Lui CY, Ornato JP, Buell JC. Lack of superiority of the vectorcardiogram over the electrocardiogram in detecting inferior wall myocardial infarction regardless of time since infarction. *J Electrocardiol* 1987;20:241–246
14. Edenbrandt L, Pahlm O, Lyttkens K. Vectorcardiogram more sensitive than 12-lead ECG in the detection of inferior myocardial infarction. *Clin Physiol* 1990;10:551–559
15. Mehta J, Hoffman I, Smedresman P. Vectorcardiographic, electrocardiographic, and angiographic correlations in apparently isolated inferior wall myocardial infarction. *Am Heart J* 1976;91:699–704
16. Tranchesi J, Moffa PJ, Pastore CA. Block of the antero-medial division of the left bundle branch of His in coronary diseases. Vectorcardiographic characterization. *Arq Bras Cardiol* 1979;32:355–360
17. Nakaya Y, Hiraga T. Reassessment of the subdivision block of the left bundle branch. *Jpn Circ J* 1981;45:503–516
18. Inoue H, Nakaya Y, Niki T, Mori H, Hiasa Y. Vectorcardiographic and epicardial activation studies on experimentally-induced subdivision block of the left bundle branch. *Jpn Circ J* 1983;47:1179–1189
19. Rosenbaum MB, Yesurun J, Lazzari JO, Elizari MV. Left anterior hemiblock obscuring the diagnosis of right bundle branch block. *Circulation* 1973;48:298–303
20. Rosenbaum MB, Elizari MV, Lazzari JO, Halpern MS, Nau GJ. Bilateral bundle branch block: its recognition and significance. *Cardiovasc Clin* 1971;2:151–179
21. Guo X, Jue X, Ruan Y. Model TJ-IV computer-assisted vectorcardiogram analysis system. *J Tongji Med Univ* 2001;21:22–81
22. Ellison RC, Restieaux NJ. *Vectorcardiography in Congenital Heart Disease. A Method for Estimating Severity*. p 44. Philadelphia–London-Toronto: W.B. Saunders Company; 1972.; Chapter 5
23. Ellison RC, Restieaux NJ. *Vectorcardiography in Congenital Heart Disease. A Method for Estimating Severity, Valvular Pulmonic Stenosis* pp 60–74. Philadelphia–London-Toronto: W.B. Saunders Company; 1972.; Chapter 6
24. Giorgi C, Nadeau R, Primeau R. Comparative accuracy of the vectorcardiogram and electrocardiogram in the localization of the accessory pathway in patients with Wolff-Parkinson-White syndrome: validation of a new vectorcardiographic algorithm by intraoperative epicardial mapping and electrophysiologic studies. *Am Heart J* 1990;119:592–598
25. Pastore CA, Moffa PJ, Tobias NM, de Moraes AP, Nishioka SA, et al. Segmental blocks of the right bundle-branch and electrically inactive areas. Differential electro-vectorcardiographic diagnosis. *Arq Bras Cardiol* 1985;45:309–317
26. Pastore CA, Moffa PJ, Spiritus MO. Fascicular blocks of the right branch. Standardization of vectorelectrocardiographic findings. *Arq Bras Cardiol* 1983;41:161–166
27. Atarashi H, Ogawa S, Harumi K. Idiopathic Ventricular Fibrillation Investigators. Three-year follow-up of patients with right bundle branch block and ST segment elevation in the right precordial leads: Japanese Registry of Brugada Syndrome. Idiopathic Ventricular Fibrillation Investigators. *J Am Coll Cardiol* 2001;37:1916–1920
28. Pérez Riera AR, Ferreira C, Schapachnik E. Value of 12 lead electrocardiogram and //derived methodologies in the diagnosis of Brugada disease”. In *The Brugada Syndrome From Bench to Bedside*. (Eds. Antzelevich C, Brugada P, Brugada J, Brugada R),pp 87–110. Futura: Blackwell; 2005.; Chapter 7
29. Riera AR, de Cano SJ, Cano MN. Vector electrocardiographic alterations after percutaneous septal ablation in obstructive hypertrophic cardiomyopathy. Possible anatomic causes. *Arq Bras Cardiol* 2002;79:466–475
30. Guo XM, Que XH, Ma YX, Wang ZC. Development and applications of an auto-analyzing system for model TJ-IV vector-cardiogram. *Zhongguo Yi Liao Qi Xie Za Zhi* 2005;29(1):19–22
31. Rautaharju PM. A hundred years of progress in electrocardiography. 2: The rise and decline of vectorcardiography. *Can J Cardiol* 1998;4:60–71
32. Grishman A, Donoso E. Spatial vectorcardiography II. *Mod Concepts Cardiovasc Dis* 1961;30:693–696
33. Benchimol A, Desser KB, Schumacher J. Left anterior hemiblock from inferior infarction with left axis deviation. *Chest* 1972;61:74–76