

Supplementary table 1. Outcomes by definitive nonoperative management (NOM) strategy and blunt splenic injury grade

Outcome, n (%)	<u>Grades I and II</u>		<u>Grade III</u>		<u>Grade IV</u>		<u>Grade V</u>	
	Observation (n=201)	SAE (n=14)	Observation (n=92)	SAE (n=34)	Observation (n=30)	SAE (n=33)	Observation (n=2)	SAE (n=11)
Mortality	4 (2.0)	2 (14.3)*	4 (4.4)	9 (0)	1 (3.3)	1 (3.0)	0 (0)	0 (0)
ICU admission	152 (75.6)	14 (100)*	78 (84.8)	33 (97.1)*	27 (90.0)	33 (100)*	2 (100)	10 (90.9)
# blood products ^a	5 (3-12)	1.5 (1-3)*	2 (2-6)	2 (2-6)	2.5 (1-8)	4.5 (3-6)	1 (1-1)	5 (2-9.5)*
Hospital LOS ^a	6 (3-11)	9 (5-16)**	5 (3-10)	11 (4-14)**	5 (4-7)	8 (5-14)*	5 (4-6)	7 (4-11)
ICU LOS ^a	2 (1-5)	7.5 (2-15)**	2 (1-4)	4 (2-7)**	3 (1.5-4.5)	4 (3-9)**	2.5 (2-3)	4 (2-6)
Failed NOM	0 (0)	0 (0)	4 (4.4)	0 (0)	0 (0)	1 (3.0)	0 (0)	0 (0)

SAE, splenic artery embolization; ICU, intensive care unit.

^aData presented as median, interquartile range

*P < 0.20; ** P < 0.05.