

Supporting Information for

Original Article

Matrine attenuates oxidative stress and cardiomyocyte apoptosis in doxorubicin-induced cardiotoxicity *via* maintaining AMPK α /UCP2 pathway

Can Hu^{a,b,c,†}, Xin Zhang^{a,b,c,†}, Wenyong Wei^{a,b,c}, Ning Zhang^{a,b,c}, Haiming Wu^{a,b,c},
Zhenguo Ma^{a,b,c}, Lingli Li^{a,b,c}, Wei Deng^{a,b,c,*}, Qizhu Tang^{a,b,c,*}

^aDepartment of Cardiology, Renmin Hospital of Wuhan University, Wuhan 430060, China

^bCardiovascular Research Institute of Wuhan University, Wuhan 430060, China

^cHubei Key Laboratory of Cardiology, Wuhan 430060, China

Received 11 January 2019; received in revised form 2 February 2019; accepted 11 March 2019

*Corresponding authors. Tel.: +86 27 88073385; fax: +86 27 88042292.

E-mail addresses: qztang@whu.edu.cn (Qizhu Tang), vivideng1982@whu.edu.cn (Wei Deng).

†These authors made equal contributions to this work.




Figure S1 Efficiency of shAmpka2 in H9C2 cells. (A) Western blot for AMPK α and ACC ($n=6$). (B) Quantitative data for AMPK α and ACC ($n=6$). Values represent the mean \pm SEM.

* $P<0.05$ versus the matched groups.