

Nano-hydroxyapatite in oral care cosmetics: characterization and cytotoxicity assessment

Catarina C. Coelho^{a,b,c,d,*}, Liliana Grenho^{e,f}, Pedro S. Gomes^{e,f}, Paulo A. Quadros^d,
Maria H. Fernandes^{e,f}

^a*Instituto de Investigação e Inovação em Saúde, U.Porto, Porto, Portugal*

Address: Rua Alfredo Allen, 208, 4200-135 Porto, Portugal

^b*Instituto de Engenharia Biomédica, U.Porto, Porto, Portugal*

Address: Rua Alfredo Allen, 208, 4200-135 Porto, Portugal

^c*Faculdade de Engenharia, U.Porto, Porto, Portugal*

Address: Rua Dr. Roberto Frias, s/n, 4200-465 Porto, Portugal

^d*FLUIDINOVA, S.A., Maia, Portugal*

*Address: Rua Engenheiro Frederico Ulrich, 2650, 4470-605 Moreira da Maia,
Portugal*


^e*Laboratory for Bone Metabolism and Regeneration, Faculdade de Medicina Dentária,
U.Porto, Porto, Portugal*

Address: Rua Dr. Manuel Pereira da Silva, s/n, 4200-393 Porto, Portugal


^f*LAQV/REQUIMTE, U.Porto, Porto, 4160-007, Portugal*

**Corresponding author (catarina.coelho@i3s.up.pt)*


Supplementary Information



Supplementary Figure 1: TEM images of the hydroxyapatite nanoparticles before and after sterilization using autoclave.



Supplementary Figure 2: FTIR spectra for hydroxyapatite nanoparticles before and after sterilization by autoclave.



Supplementary Figure 3: XRD patterns obtained for hydroxyapatite nanoparticles before and after sterilization with autoclave.