


Appendix 3

Network meta-analysis of eHealth modalities used to deliver interventions for the reduction of pain severity in a chronic pain population.



Pairwise comparisons between eHealth modalities delivering interventions for a reduction in pain severity

Modality	A	B	C	D	E	F	G
B	-0.07 [-.67, .65]						
C	-.26 [-1.04, .51]	-.26 [-1.24, .73]					
D	.15 [-.67, .96]	.16 [-.88, 1.2]	.41 [-.71, 1.53]				
E	-.06 [-.97, .85]	-.05 [-1.15, 1.05]	.2 [-.95, 1.36]	-.21 [-1.43, 1.02]			
F	2.22 [1.35, 3.08]	2.22 [1.18, 3.27]	2.48 [1.33, 3.63]	2.07 [.88, 3.26]	2.28 [1.02, 3.54]		
G	.309 [.02, .6]	0.32 [-.33, .98]	.57 [-.25, 1.4]	.16 [-.7, 1.03]	0.37 [-.58, 1.33]	-1.91 [-2.72, -1.1]	

H	.2 [-.003, .41]	.21 [-.44, .87]	.047 [-.27, 1.21]	.05 [-.79, .9]	.27 [-.62, 1.15]	-2.01 [-2.9, -1.13]	-.11 [-.46, .25]
----------	---------------------------	--------------------	----------------------	-------------------	---------------------	-------------------------------	---------------------

Data displayed as SMD [Credible Interval]

Bolded entries are statistically significant

A=Internet, B=Virtual reality, C=Telephone, D=Mobile applications, E=Interactive voice response, F=Video teleconferencing, G=Enhanced control, H=Control
