**Title:** Expression and secretion of CXCL12 are enhanced in autosomal dominant polycystic kidney disease

**Author's name:** Hyunho Kim<sup>1</sup>, Jinmo Sung<sup>1</sup>, Hyunsuk Kim<sup>2</sup>, Hyunjin Ryu<sup>3</sup>, Hayne Cho Park<sup>4</sup>, Yun Kyu Oh<sup>5</sup>, Hyun-seob Lee<sup>6</sup>, Kook-Hwan Oh<sup>3</sup>, Curie Ahn<sup>3,\*</sup>

**Affiliation:** <sup>1</sup> Center for Medical Innovation, Biomedical Research Institute, Seoul National University Hospital, Seoul, Republic of Korea

<sup>2</sup> Internal Medicine, Hallym University Medical Center, Chuncheon Sacred Heart Hospital, Chuncheon-si, Gangwon-do, Republic of Korea

<sup>3</sup> Department of Internal Medicine, Seoul National University College of Medicine, Seoul, Republic of Korea

<sup>4</sup> Department of Internal Medicine, Hallym University Medical Center, Kangnam Sacred Heart Hospital, Seoul, Republic of Korea

<sup>5</sup> Department of Internal Medicine, Seoul National University, Boramae Medical Center, Seoul, Republic of Korea


<sup>6</sup> Genomics Core Facility, Department of Transdisciplinary Research and Collaboration, Biomedical Research Institute, Seoul National University Hospital, Seoul, Republic of Korea

Running Title: Upregulation of CXCL12 in renal cyst cells

**Keywords:** Autosomal dominant polycystic kidney disease (ADPKD), Cyst, hTERT, CXCL12, CXCR4.

\*Corresponding Author's Information: Tel: +82-2-2072-2222; Fax: +82-2-741-4876; E-mail: curie@snu.ac.kr

## **Supplementary Figure**



**Figure S1. CXCL12 protein-protein interaction network analysis.** The functional interaction network among 42 differentially expressed genes of the significant GO term of upregulated genes in the ADPKD renal cyst cell lines (RC1, RC2, RC3) vs the normal proximal tubular cell line (RPTEC/TERT1) using STRING v11.0.