

SUPPLEMENTARY MATERIALS

(Incidence of Ductal Carcinoma in Situ in the United States, 2000-2014, Ryser et al.)

ICD-0-3 histology codes included in the study:

8050/2: Papillary carcinoma in situ
8140/2: Adenocarcinoma in situ, NOS
8200/2: Adenoid cystic carcinoma in situ
8201/2: Cribriform carcinoma in situ
8211/2: Tubular adenocarcinoma in situ
8230/2: Ductal carcinoma in situ, solid type
8246/2: Neuroendocrine carcinoma in situ
8260/2: Papillary adenocarcinoma in situ, NOS
8343/2: Papillary carcinoma, encapsulated, in situ
8401/2: Apocrine adenocarcinoma in situ
8480/2: Mucinous adenocarcinoma in situ
8481/2: Mucin-producing adenocarcinoma in situ
8490/2: Signet ring cell carcinoma in situ
8500/2: Intraductal carcinoma, noninfiltrating, NOS
8501/2: Comedocarcinoma, noninfiltrating
8503/2: Noninfiltrating intraductal papillary adenocarcinoma
8504/2: Noninfiltrating intracystic carcinoma
8507/2: Intraductal micropapillary carcinoma
8508/2: Cystic hypersecretory carcinoma, in situ
8521/2: Ductular carcinoma in situ
8522/2: Intraductal and lobular in situ carcinoma
8523/2: Intraductal with other types of carcinoma in situ
8543/2: Paget disease in situ and intraductal carcinoma
8550/2: Acinar cell carcinoma in situ