

PROTOCOL DEVIATIONS

Table 11: Off-Protocol Events by Site

Off-Protocol Events	Gulu (N=379) N(%)	Kitgum (N=375) N(%)	Pader (N=386) N(%)	Total (N=1140)
Participants enrolled outside the target period	8(2.11)	10(2.67)	5(1.30)	23(2.02)
Number of health centers that did not meet the target enrolment (N=30)	8(80)	5(50)	3(30)	16(53.33)
Enrolled participants did not meet criteria for major depression	1(7.69)	7(53.85)	5(38.46)	13(1.14)
Enrolled participants with high suicide risk	4(12.50)	5(15.63)	23(71.88)	32(2.81)
Did not have baseline assessments	12(57.14)	4(19.05)	5(23.81)	21(1.84)
Lost to follow up after end of treatment	37(57.81)	11(17.19)	16(25)	64(5.61)
Group session attendance of trial participants				
Did not attend any group session	28(7.39)	12(3.20)	8(2.07)	48(4.21)
Attended one group session	3(0.79)	1(0.27)	1(0.26)	5(0.44)
Attended two group sessions	0(0.00)	2(0.53)	3(0.78)	5(0.44)
Attended three group sessions	3(0.79)	1(0.27)	1(0.26)	5(0.44)
Attended four group sessions	7(1.85)	0(0.00)	7(1.81)	14(1.23)
Attended five group sessions	5(1.32)	3(0.80)	6(1.55)	14(1.23)
Attended six group sessions	11(2.90)	11(2.93)	19(4.92)	41(3.60)
Attended seven group sessions	22(5.80)	19(5.07)	15(3.89)	56(4.91)
Attended eight group sessions	300(79.16)	326(86.93)	326(84.46)	952(83.51)