


Supplemental Table 1. Association between race and enrollment by region or poverty
Of note, some of the subgroups in these post-hoc analyses contain small numbers of patients and should thus be interpreted with caution.

	N	Rate of Enrollment (%)	Crude Risk Difference (%)	Adjusted* Risk Ratio
Stratified by Region				
Northeast	50			
White, non Hispanic	33	73	ref	ref
Black, non Hispanic	10	40	-33	0.46 (0.36, 0.58)
Hispanic	2	100	27	1.34 (1.06, 1.68)
Other	5	20	-53	0.17 (0.07, 0.43)
Midwest	98			
White, non Hispanic	70	57	ref	ref
Black, non Hispanic	12	17	-40	0.50 (0.29, 0.88)
Hispanic	6	50	-7	0.94 (0.89, 1.53)
Other	10	50	-7	0.97 (0.63, 1.50)
South	124			
White, non Hispanic	54	57	ref	ref
Black, non Hispanic	23	30	-27	0.44 (0.23, 0.84)
Hispanic	27	52	-5	0.94 (0.66, 1.36)
Other	20	50	-7	0.66 (0.36, 1.22)
West	98			
White, non Hispanic	41	63	ref	ref
Black, non Hispanic	3	67	4	1.02 (0.32, 3.28)
Hispanic	37	65	2	1.01 (0.63, 1.60)
Other	17	65	2	1.06 (0.84, 1.33)
Stratified by Zip-based Poverty				
>4% poverty	281			
White, non Hispanic	137	65	ref	ref
Black	43	30	-35	0.49 (0.34, 0.70)
Hispanic	62	65	0	0.94 (0.77, 1.15)
Other	39	59	-6	0.87 (0.66, 1.14)
<4% poverty	81			
White, non Hispanic	58	52	ref	ref
Black	4	50	-2	1.16 (0.81, 1.66)
Hispanic	8	25	-27	0.41 (0.11, 1.51)
Other	11	27	-25	0.42 (0.17, 1.08)

*Adjusted for race, age, year, SES, and acuity

Supplemental Figure 1. Institution-level variation in the rates of trial enrollment on AAML1031 (left) and 9-month inpatient mortality (right) Institutions are numbered in order of decreasing rates of trial enrollment. Enrollment and mortality do not appear to be correlated on the institution-level.



Supplemental Table 2. Institution-level characteristics as predictors of enrollment
None of the institutional characteristics were associated with trial enrollment.

Characteristics	Unadjusted RR (95% CI)	p
AML patient volume (by 50 patients)	1.17 (0.28, 4.94)	0.83
% Black*	0.94 (0.85, 1.04)	0.23
% public insurance*	1.04 (0.97, 1.10)	0.27
% urban*	1.01 (0.92, 1.10)	0.88
% hospital-wide inpatient mortality (1% increase)	0.76 (0.47, 1.24)	0.27
Median Household income (by 10k)	1.05 (0.91, 1.21)	0.53

* 10% increase

Supplemental Table 3. Characteristics of matched versus unmatched patients

Those who were unable to be matched did not differ significantly from those that were able to be matched.

	AAML1031 COG-PHIS merged patients matched to PHIS AML cohort N=206 (%)	AAML1031 COG- PHIS merged patients unmatched to PHIS AML cohort N=39 (%)	p-value
Race/Ethnicity			0.84
White Non-Hispanic	119 (58)	24 (62)	
Black Non-Hispanic	21 (10)	5 (13)	
Hispanic	42 (20)	7 (18)	
Other	24 (12)	3 (8)	
Sex			0.87
Female	98 (48)	18 (46)	
Male	108 (52)	21 (54)	
Age, years			0.11
0-1	15 (7)	1 (3)	
1-5	54 (26)	9 (23)	
5-10	37 (18)	5 (13)	
10-15	52 (25)	7 (18)	
>15	48 (23)	17 (44)	
Insurance type			0.28
Private	100 (49)	24 (62)	
Public	97 (47)	13 (33)	
Other	9 (4)	2 (5)	
Illness severity at Presentation			0.25
No ICU	187 (91)	38 (97)	
1 system	14 (7)	0 (0)	
2 or more systems	5 (2)	1 (3)	
Region			0.26
South	62 (30)	10 (26)	
Midwest	50 (24)	15 (39)	
Northeast	31 (15)	3 (8)	
West	63 (31)	11 (28)	
Year			0.77
2011	21 (10)	2 (5)	
2012	82 (40)	16 (41)	
2013	68 (33)	13 (33)	
2014	35 (17)	8 (21)	