

Supplement to

“Bacteria isolated from the cuticle of plant-parasitic nematodes attached to and antagonized the root-knot nematode *Meloidogyne hapla*”

Olivera Topalović¹, Ahmed Elhady^{1,2}, Johannes Hallmann¹, Katja R. Richert-Pöggeler¹, Holger Heuer^{1*}

¹ Dept. Epidemiology and Pathogen Diagnostics, Julius Kühn-Institut — Federal Research Centre for Cultivated Plants, Braunschweig, Germany

² Department of Plant Protection, Faculty of Agriculture, Benha University, Benha, Egypt

*Correspondence: holger.heuer@julius-kuehn.de

Supplementary Table S1. Identification of bacterial strains that were highly attached to J2 of *Meloidogyne hapla* or *Pratylenchus penetrans*, using 16S rRNA gene sequencing.

Isolates	Soil source	GenBank BLAST hit	Accession ID, % identity
S10-12 (Mh)	Sickte (field 10)	<i>Acinetobacter</i> sp.	KX_982223.1, 99%
E1 (Mh)	Elsdorf	<i>Acinetobacter</i> sp.	KJ_147068.1, 99%
i.44 (Mh)	Geisenheim	<i>Microbacterium foliorum</i>	CP_019892.1, 99%
i.47 (Mh)	Geisenheim	<i>Microbacterium</i> sp.	CP_018151.1, 99%
BS1-7 (Mh)	Dahnsdorf (field BS1)	<i>Brevundimonas vesicularis</i>	MH_144292.1, 99%
S5-5 (Mh)	Quedlinburg (Schlag 5)	<i>Brevundimonas</i> sp.	KY_907035.1, 99%
G1 (Mh)	Geisenheim	<i>Kocuria uropiqioeca</i>	NR_157676.1, 99%
i.10 (Mh)	Geisenheim	<i>Microbacterium phyllosphaerae</i>	NR_025405.1, 99%
K5 (Mh)	Klein Wanzleben	<i>Micrococcus</i> sp.	MH_671525.1, 99%
G11 (Mh)	Geisenheim	<i>Sphingopyxis</i> sp.	EF_540469.1, 99%
BS1-2 (Mh)	Dahnsdorf (Schlag 2)	<i>Sphingopyxis</i> sp.	AB_161684.1, 99%
K6 (Mh)	Klein Wanzleben	<i>Microbacterium</i> sp.	MG_266365.1, 99%
K1 (Mh)	Klein Wanzleben	<i>Staphylococcus equorum</i>	MF_578770.1, 99%
i.20 (Mh)	Geisenheim	<i>Microbacterium</i> sp.	EU_747700.1, 99%
i.14 (Pp)	Bundesallee (bulk soil)	<i>Microbacterium oxydans</i>	MF_459692.1, 100%
i.27 (Pp)	Bundesallee (maize rhizosphere)	<i>Pseudomonas protegens</i>	NR_114749.1, 100%
i.37 (Pp)	Bundesallee (soybean rhizosphere)	<i>Alcaligenes faecalis</i>	NR_113606.1, 100%