

1 **Supplementary information**

2 A co-expressed gene status of adenylate kinase 1/4 reveals prognostic gene
3 signature associated with prognosis and sensitivity to EGFR targeted therapy
4 in lung adenocarcinoma

5

6 Yi-Hua Jan^{1*}, Tsung-Ching Lai^{1*}, Chih-Jen Yang², Ming-Shyan Huang^{4#}, and
7 Michael Hsiao^{1, 3#}

8

9 ¹Genomics Research Center, Academia Sinica, Taipei, Taiwan, ² Department
10 of Internal Medicine, Kaohsiung Medical University Hospital, School of
11 Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan, ³Department of
12 Biochemistry, College of Medicine, Kaohsiung Medical University, Kaohsiung,
13 Taiwan. ⁴School of Medicine, I-Shou University, Kaohsiung, Taiwan.

14

15 *The first two authors contribute equally to this works.

16 #The last two authors are senior authors and contribute equally to this works.

17

18 Correspondence to: Dr. Michael Hsiao, Genomics Research Center, Academic
19 Sinica, 128 Academia Road, Section 2, Taipei 115, Taiwan. Phone: 886-2-
20 2787-1243; Fax: 886-2-2789-9931; E-mail: mhsiao@gate.sinica.edu.tw

21

22

Figure S3.






Figure S3.





Fig 5.





Fig 5.





Fig 5.

H358-shAK4-2
H358-shAK4-1
H358-shLuc





Fig 5.




Fig 5.

beta-actin

H358-shAK4-2
H358-shAK4-1
H358-shLuc





Fig 5.





Fig 5.





Fig 5.





Fig 5.

