

ESM_2

**RELEVANCE OF VEGFA IN RAT LIVERS SUBJECTED TO PARTIAL HEPATECTOMY
UNDER ISCHEMIA-REPERFUSION
JOURNAL OF MOLECULAR MEDICINE**

Esther Bujaldon^{1*}, María Eugenia Cornide-Petronio^{1*}, José Gulfo², Floriana Rotondo¹, Cindy Ávalos de León¹, Elsa Negrete-Sánchez¹, Jordi Gracia-Sancho³, Anna Novials^{4,5}, Mónica B. Jiménez-Castro^{6#}, Carmen Peralta^{1, 2, 7#}

¹Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain; ²Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas, Barcelona, Spain. ³Liver Vascular Biology Research Group, IDIBAPS, CIBEREHD, Barcelona, Spain; ⁴Diabetes and Obesity Research Laboratory, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain; ⁵Spanish Biomedical Research Center in Diabetes and Associated Metabolic Disorders (CIBERDEM), Barcelona, Spain; ⁶Transplant Biomedicals S.L., Barcelona, Spain; ⁷Facultad de Medicina, Universidad International de Cataluña, Barcelona, Spain.

* EB and MECP contributed equally to this work (as first author)

MJC and CP contributed equally to this work (as last author)

Correspondence to: Carmen Peralta Uroz


E-mail: cperalta@clinic.ub.es

Telephone: +34932275400 Ext 4177

SUPPLEMENTARY FIGURE LEGENDS

Supplementary Figure 2. Effect of VEGFA on hepatic damage and regeneration in Ob Zucker rats undergoing PH+I/R 72 h after surgery. (A) Plasma AST, ALT, GLDH, damage score, ALP and bilirubin levels. (B) Hepatic regeneration (percentage of Ki67-positive-hepatocytes, representative photomicrographs of Ki-67 immunohistochemical positivity (10x) and cyclin E levels). +P < 0.05 versus PH+I/R.

A Hepatic damage



B Liver regeneration

