GRADE evidence profile

Certainty assessment							№ of patients		Effect		
№ of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Mindfulness	control	Relative (95% CI)	Absolute (95% CI)	Certainty
Pain End of treatment											
7	randomised trials	serious a	serious ^b	not serious	not serious	none	263	331	-	SMD 0.46 SD lower (0.75 lower to 0.17 lower)	Low
Pain Follow-up											
6	randomised trials	serious °	not serious	not serious	serious ^d	none	224	313	-	SMD 0.25 SD lower (0.52 lower to 0.01 higher)	⊕⊕⊖⊖ Low
Depressi	Depression End of treatment										
8	randomised trials	serious a	serious ^b	not serious	not serious	none	277	347	-	SMD 0.49 SD lower (0.85 lower to 0.12 lower)	⊕⊕⊖ Low
Depressi	on Follow-up										
7	randomised trials	serious °	serious ^b	not serious	not serious	none	238	329	-	SMD 0.48 SD lower (0.77 lower to 0.19 lower)	ФФО Low
Anxiety I	End of treatme	nt									
4	randomised trials	serious a	not serious	not serious	not serious	none	150	228	-	SMD 0.37 SD lower (0.71 lower to 0.02 lower)	⊕⊕⊕⊖ MODERATE
Anxiety F	Follow-up		I	1	1		ı	1		1	
3	randomised trials	serious °	serious b	not serious	serious ^d	none	111	210	-	SMD 0.44 SD lower (0.9 lower to 0.02 higher)	⊕⊖⊖ VERY LOW

Sleep quality End of treatment

3	randomised trials	serious °	not serious	not serious	serious ^d	none	126	173	-	SMD 0.33 SD lower (0.7 lower to 0.04 higher)	LOW LOW
Sleep qu	ı ality Follow-up)	I	l	I	l	1	I	I		
3	randomised trials	serious º	not serious	not serious	not serious	none	128	175	-	SMD 0.25 SD lower (0.5 lower to 0)	⊕⊕⊕○ MODERATE
Health-re	l elated quality o	of Life End of tr	l eatment							l	
2	randomised trials	serious e	serious ^b	not serious	serious ^d	none	92	152	-	SMD 0.62 SD lower (1.52 lower to 0.28 higher)	⊕⊖⊖ VERY LOW
Health-re	Health-related quality of Life Follow-up										
2	randomised trials	serious e	serious ^b	not serious	serious ^d	none	92	152	-	SMD 0.62 SD lower (1.52 lower to 0.28 higher)	⊕⊖⊖ VERY LOW
Mindfuln	ess End of tre	atment									
2	randomised trials	serious e	not serious	not serious	not serious	none	75	134	-	SMD 0.4 SD lower (0.69 lower to 0.11 lower)	⊕⊕⊕⊖ MODERATE
Mindfuln	ess Follow-up										
2	randomised trials	serious °	not serious	not serious	serious ^d	none	77	136	-	SMD 0.28 SD lower (0.56 lower to 0.01 higher)	⊕⊕⊖⊖ LOW
Fatigue -	Fatigue - not measured										
									-	0 (0 to 0)	-
Workabil	Workability - not measured										
									-	0 (0 to 0)	-
					<u> </u>						

Explanations

- a. The Risk of Bias show High Risk on attrition bias, selection bias and performance bias. In addition there are some uncertainty and unclear risk.
- b. The p-value is low and the I-square is high.
- c. The Risk of Bias show High Risk on attrition bias and performance bias. In addition there are some uncertainty and unclear risk.
- d. The confidence interval goes from a large effect in favor of mindfulness to a small effect in favor of control.
- e. The Risk of Bias show High Risk on performance bias. In addition there are some uncertainty and unclear risk.