

Supplementary Information for

The Effect of Dietary Supplementation of Crocetin for Myopia Control in Children: A Randomized Clinical Trial

Kiwako Mori^{1,2,4}, Hidemasa Torii^{1,2,4}, Satoko Fujimoto^{3,4}, Xiaoyan Jiang^{1,2}, Shin-ichi Ikeda^{1,2}, Erisa Yotsukura^{1,2}, Shizuka Koh³, Toshihide Kurihara^{1,2,*}, Kohji Nishida^{3,*}, Kazuo Tsubota^{1,*}

Supplementary Table 1

Table S1	Safety					
Subject	69 patients (male:38 patients, female:31 patients)					
Adverse reactions	0 patient 0 case					
Adverse events	30 patients 50 cases (All: Not Related)					
	Influenza A infection	2	Cold	9	Fracture (Left ankle)	1
	Influenza B infection	8	Itchy eyes	1	Parotitis	1
	EB virus infection	1	Bronchitis	1	Headache	1
	Allergic conjunctivitis	2	Seasonal allergic rhinitis	1	Fever	2
	Virus conjunctivitis (left eye)	1	Acute conjunctivitis	1	Sprain (right big toe)	1
	Osteochondrosis of tarsus	1	Acute conjunctivitis (left eye)	1	Streptococcal infection	2
	Osteochondrosis of tarsus (left heel)	1	Acute upper respiratory infection	1	Nausea	1
	Gastroenteritis	1	Acute sinusitis	1	Cervical lymphadenopathy	1
	Sore throat	2	Acute chalazion (left eye)	1	hordeolum (right eye)	1
	Infectious gastroenteritis	1	Stomatitis	2		
Withdrawn	2 patients. Less intake (<i>n</i> = 1), inconvenient (<i>n</i> = 1)					