Association between Galectin-3 levels within central and peripheral venous blood, and adverse left ventricular remodelling after first acute myocardial infarction

Olivera M. Andrejić^a, Rada M. Vučić^{b,c}, Milan Pavlović^{d,i}, Lana McClements^f, Dragana Stokanović^g, Tatjana Jevtović –Stoimenov^h, Valentina N. Nikolić^g

^aClinic for Pulmonary Diseases, Clinical Centre Kragujevac, Zmaj Jovina Street 30, 34000 Kragujevac, Serbia

^bUniversity of Kragujevac, Serbia, Faculty of Medical Sciences, Department of Internal medicine, Svetozara Markovica Street 69, 34000 Kragujevac, Serbia

^cClinic for Cardiovascular Diseases, Clinical Centre Kragujevac, Zmaj Jovina Street 30, 34000 Kragujevac, Serbia

^dDepartment of Internal Medicine - Cardiology, Medical Faculty, University of Nis, Bulevar dr Zorana Djindjica 81, Nis, Serbia

ⁱClinic for Cardiovascular Diseases, Clinical Centre Nis, Bulevar dr Zorana Djindjica 48, Nis, Serbia

^fSchool of Life Sciences, Faculty of Science, University of Technology Sydney, Sydney, PO Box 123 Broadway, NSW 2007, Australia

^gDepartment of Pharmacology and Toxicology, Medical Faculty, University of Nis, Bulevar dr Zorana Djindjica 81, Nis, Serbia

^hInstitute of Biochemistry, Medical Faculty, University of Nis, Bulevar dr Zorana Djindjica 81, Nis, Serbia

Declarations of interest: None.

Financial support: This study funded by Faculty of Medicine, University of Nis, (Grant No INT-MF19), as well as Serbian Ministry of Education, Science and Techno-logical Development (Grant No III41018 and 44044).

***Reprints and correspondence**: Dr Rada M. Vučić; 00381631123610, email: rada.vucic@gmail.com; Milovana Gusica 8/15, 34000 Kragujevac, Serbia

Supplementary Figure 2. Schematic of the study events.

