

SUPPLEMENTARY TABLE S6. ASSOCIATIONS OF PRE- AND EARLY PREGNANCY DEPRESSIVE SYMPTOMS WITH GESTATIONAL WEIGHT GAIN RATE (LBS/WEEK) AMONG WOMEN WITH MEASURED PREPREGNANCY WEIGHT

	N	Mean difference ^a (95% CI)	N	RR ^a (95% CI) for GWG below the IOM recommendations	N	RR ^a (95% CI) for GWG above the IOM recommendations
No depressive symptoms in pre- or early pregnancy	62,807	Reference	14,798	Reference	34,353	Reference
Prepregnancy onset depression	7357	0.00 (-0.01 to 0.02)	1793	1.11 (1.07 to 1.15)	4250	1.03 (1.02 to 1.05)
Early pregnancy onset depression	6549	0.04 (0.02 to 0.07)	1536	1.04 (0.98 to 1.11)	3728	1.05 (1.02 to 1.08)

^aModel is adjusted for maternal age, race/ethnicity (non-Hispanic white, non-Hispanic black, Hispanic, Asian/Pacific Islander, Native American/multiracial, other/unknown), nulliparity (Y/N), Medicaid (Y/N), smoking in first 20 weeks of pregnancy (Y/N), alcohol use in first 20 weeks of pregnancy (Y/N), prepregnancy BMI category (underweight, normal weight, overweight, obese), and severity of early pregnancy depressive symptoms (mild, moderate, severe).