

Supplementary table S1: additional treatment with avastin stratified by (a) TCGA subtype and (b) EGFR amplification status

a

	Classical	Mesenchymal	Proneural	p Value
	n (%)	n (%)	n (%)	
<u>Number of treatment pre bevacizumab</u>				
0	1 (5%)	1 (5%)	0 (0%)	0.453
1	5 (26%)	5 (24%)	5 (22%)	
2	7 (37%)	9 (43%)	10 (43%)	
3	4 (21%)	6 (29%)	3 (13%)	
4	2 (11%)	0 (0%)	5 (22%)	
<u>Number of treatment during bevacizumab</u>				
0	11 (58%)	9 (43%)	9 (39%)	0.588
1	5 (26%)	10 (48%)	10 (43%)	
2	2 (11%)	2 (10%)	4 (17%)	
3	1 (5%)	0 (0%)	0 (0%)	
<u>Number of treatment post bevacizumab</u>				
0	10 (53%)	13 (62%)	17 (74%)	0.133
1	5 (26%)	7 (33%)	3 (13%)	
2	4 (21%)	0 (0%)	1 (4%)	
3	0 (0%)	1 (5%)	1 (4%)	
4	0 (0%)	0 (0%)	1 (4%)	

b

EGFR amplification status				
	No	Yes	Unknown	p value
<u>Number of treatment pre bevacizumab</u>				
0	0 (0%)	2 (9%)	0 (0%)	0,298
1	8 (30%)	6 (26%)	1 (8%)	
2	10 (37%)	7 (30%)	9 (69%)	
3	5 (19%)	5 (22%)	3 (23%)	
4	4 (15%)	3 (13%)	0 (0%)	
<u>Number of treatment during bevacizumab</u>				
0	11 (41%)	11 (48%)	7 (54%)	0,785
1	13 (48%)	7 (30%)	5 (38%)	
2	3 (11%)	4 (17%)	1 (8%)	
3	0 (0%)	1 (4%)	0 (0%)	
<u>Number of treatment post bevacizumab</u>				
0	20 (74%)	15 (65%)	5 (38%)	0,109
1	5 (19%)	5 (22%)	5 (38%)	
2	1 (4%)	3 (13%)	1 (8%)	
3	0 (0%)	0 (0%)	2 (15%)	
4	1 (4%)	0 (0%)	0 (0%)	