

Appendix Table 4: Bootstrapping, long-term opioid use

| Initial Provider | N | OR (95% CI) bootstrapping | Long-term use, OR (95% CI) |
|--------------------|-----|------------------------------|-------------------------------|
| Physical therapy | 200 | 0.26 (0.16, 0.50) | 0.27 (0.15,0.48) |
| Chiropractor | 200 | 0.22 (0.18, 0.26) | 0.22 (0.18,0.26) |
| Acupuncture | 200 | | 0.07 (0.01,0.48) |
| Orthopedics | 200 | 1.09 (0.92, 1.31) | 1.10 (0.92,1.30) |
| Emergency Medicine | 200 | 0.92 (0.77, 1.11) | 0.92 (0.77,1.10) |
| Neurosurgeon | 200 | 1.45 (0.88, 2.75) | 1.50 (0.88,2.58) |
| MD Other | 200 | 2.02 (1.71, 2.41) | 2.03 (1.70,2.41) |
| Radiology | 200 | 1.22 (1.08, 1.38) | 1.22 (1.08,1.39) |
| Rehab | 200 | 1.77 (1.42, 2.26) | 1.78 (1.40,2.26) |
| Other | 200 | 0.88 (0.69, 1.14) | 0.88 (0.70,1.11) |

Appendix Table 4 Legend: Sample size for acupuncture was too small to report for bootstrapping.