

Supplementary

Effect of diosmin administration in patients with chronic venous disorders on selected factors affecting angiogenesis

Marcin Feldo^{1,*}, Magdalena Wójciak-Kosior², Ireneusz Sowa², Janusz Kocki³, Jacek Bogucki³, Tomasz Zubilewicz¹, Jan Kęsik¹, Anna Bogucka-Kocka⁴


¹ Department of Vascular Surgery and Angiology, Medical University of Lublin, Staszica 11, 20-081 Lublin, Poland; martinf@interia.pl (M.F.), tomasz.zubilewicz@umlub.pl (T.Z.), jankesik@umlub.pl (J.K.)

² Department of Analytical Chemistry, Medical University of Lublin, Chodźki 4a, 20-093 Lublin, Poland; kosiorma@wp.pl (M.W.K); i.sowa@umlub.pl (I.S)

³ Department of Clinical Genetics, Medical University of Lublin, Lublin, Poland, Radziwiłłowska 11, 20-080 Lublin; janusz.kocki@umlub.pl (J.K.); jacekbogucki@wp.pl (J.B.)

⁴ Chair and Department of Biology and Genetics, Medical University of Lublin, W. Chodźki 4A, 20-093 Lublin, Poland, anna.bogucka-kocka@umlub.pl


* Correspondence: martinf@interia.pl; Tel.: +48 81532 5707



(a)

(b)

Figure S1. Echo-Doppler assay obtained in patients without (a) and with (b) blood reflux.



(a)

(b)

Figure S2. Ultrasound images of patients' veins without (a) and with thrombosis marked in red (b).