

Supplementary Data

Supplementary Table S1. Model for Predicting Missing Low-Density Lipoprotein Data

	Logit model with HbA1C	Logit Model without HbA1C
Year (per year)	1.09 (1.09, 1.09)	1.09 (1.09, 1.10)
Age (per year)	1.00 (1.00, 1.00)	1.00 (1.00, 1.00)
Elixhauser comorbidities ^a	1.01 (0.99, 1.02)	1.02 (1.02, 1.02)
Primary care visits (per annual visit)	0.99 (0.99, 0.99)	0.99 (0.99, 0.99)
Race/ethnicity: non-Hispanic black	1.01 (0.99, 1.02)	1.00 (0.98, 1.02)
Race/ethnicity: Hispanic	1.00 (0.97, 1.02)	0.98 (0.96, 1.01)
Race/ethnicity: other	1.02 (1.00, 1.03)	0.90 (0.86, 0.92)
Dual use status: 50–80% VHA	1.00 (0.99, 1.01)	1.00 (0.99, 1.01)
Dual use status: 50% VHA	0.99 (0.97, 1.00)	1.00 (0.98, 1.01)
Sex: female	0.94 (0.90, 0.99)	0.93 (0.89, 0.98)
Marital status: unmarried	1.07 (1.06, 1.08)	1.07 (1.06, 1.08)
Disability >50%	0.90 (0.89, 0.92)	0.90 (0.89, 0.91)
Smoking status = yes	0.95 (0.94, 0.97)	0.95 (0.94, 0.97)
Statins prescribed: intense vs. regular	0.02 (0.02, 0.02)	0.02 (0.02, 0.02)
HbA1C >8%	1.15 (1.14, 1.17)	—
Acute coronary syndrome	1.04 (1.03, 1.06)	1.05 (1.03, 1.06)
Atherosclerotic cerebrovascular disease	1.04 (1.03, 1.06)	1.04 (1.03, 1.06)
Coronary heart disease	1.18 (1.16, 1.20)	1.18 (1.17, 1.20)
Peripheral artery disease	1.05 (1.04, 1.06)	1.05 (1.04, 1.06)

^aPer additional comorbidity.

VHA, Veterans Health Administration.

