


Supplemental Materials.


Supplemental Figure 1



Supplemental Fig. 1. Importin α 5-selective NTM suppresses lung endothelial injury (VCAM-1), decreases circulating neutrophils, and prevents apoptosis (Casp-3) in mice comprising 2 models of Endotoxin Shock: High Dose LPS model (A) and Low Dose LPS model primed with D-galactosamine (B).

Representative images (40x magnification) of lung sections in unchallenged mice (mock control) or mice challenged with LPS (**A**: 35 mg/kg LPS; **B**: 1 g/kg D-Gal then 50 μ g/kg LPS). Mice were treated with either 7 (A) or 5 (B) doses of NTM (33 mg/kg) or saline (See text for details). Data presented in this figure denotes 2 independent in vivo experiments completed with 5 mice per condition group.

Supplemental Figure. 2



Supplemental Fig. 2. Importin α 5-selective NTM reduces nuclear translocation of NF- κ B p65 (Rel A), phospho-NF- κ B p65 (p-Rel A) and phospho-STAT1 (α and β) in two models of Endotoxin Shock. A. and B. Representative immunoblots of liver nuclear extracts. Liver samples were collected 12 hr after mice were challenged with a High Dose LPS (A) or 6 hr after mice primed with D-Gal were challenged with a Low Dose LPS (B). Mice were treated with either 7 (A) or 5 (B) doses of NTM (33 mg/kg) or saline. C. and D. Quantitative representation of immunoblots shown in panel A. E. Quantitative representation of immunoblots shown in panel B. Data presented in this figure denotes 2 independent *in vivo* experiments completed with 5 mice per condition group. All signals were normalized to TATA Binding Protein (TBP) and expressed as percent inhibition \pm SEM. Significance was determined by one-way ANOVA, * - $p < 0.05$, ** - $p < 0.005$.