

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Impact of Long-Term Exposure to Wind Turbine Noise on Redemption of Sleep Medication and Antidepressants: A Nationwide Cohort Study

Aslak Harbo Poulsen, Ole Raaschou-Nielsen, Alfredo Peña, Andrea N. Hahmann, Rikke Baastrup Nordsborg, Matthias Ketzel, Jørgen Brandt, and Mette Sørensen

Table of Contents

Table S1. Characteristics of the populations for study of redemption of sleep medication and antidepressants, respectively, at start of follow-up according to residential low frequency exposure to indoor wind turbine noise calculated as mean exposure during the preceding year.

Table S2. Characteristics of the populations for study of redemption of sleep medication and antidepressants, respectively, at start of follow-up, according to residential exposure to outdoor and indoor low frequency wind turbine noise calculated as mean exposure during the preceding year.

Table S3. Pearson correlation coefficients between different noise metrics above the reference level (24 dB outdoor WTN and 5 dB indoor LF WTN) preceding entry into the study population.

Table S4. Associations between 5-year exposure to outdoor wind turbine noise and redemption of sleep medication and antidepressants in different sub-populations.

Table S5. Associations between 5-year exposure to indoor low frequency wind turbine noise and redemption of sleep medication and antidepressants in different sub-populations.